

FALLOW ROTATION WITH BARLEY AND WHEAT (POSTHARVEST - COLORADO, KANSAS, MONTANA, NEBRASKA AND WYOMING)

General Information

GENERAL INFORMATION

SOIL TEXTURE: As used on the label, “Coarse soils” are sand, loamy sand or sandy loam soils. “Medium soils” are loam, silt loam, silt, sandy clay, or sandy clay loam. “Fine soils” are silty clay, silty clay loam, clay, or clay loam. Silty clay loam soils are transitional soils and may be classified as medium textured soils in some regions of the U.S.

GENERAL PRECAUTIONS AND RESTRICTIONS

Do not allow sprays to drift on to adjacent desirable plants.

Apply this product only as specified on the label.

Do not use on other crops grown for food or forage. Observe all cautions and limitations on labeling of all products used in mixtures.

Do not rotate any crop not listed on the label for 18 months following application of DIMETRIC DF 75%.

For all uses: Low-pressure and high-volume hand-wand equipment is prohibited.

GROUND APPLICATION: Apply the proper rate of DIMETRIC DF 75% in a minimum of 10 to 40 gallons of spray mixture per acre broadcast.

Banded Application: Use proportionally less DIMETRIC DF 75% per acre in a band versus a broadcast application. For band application use 1/4 to 1 gallon of spray mix per inch of band width regardless of row spacing.

AERIAL APPLICATION: Where permitted, apply specified rate in a minimum of 2 to 10 gallons of spray mixture per acre. Do not apply aerially when wind speed is greater than 10 mph.

NOTE: Do not apply aerially when DIMETRIC DF 75% is tank-mixed with Lasso.

For All Applications of DIMETRIC DF 75%: Sprayer must be accurately calibrated before applying METRICDF. Check sprayer during application to be sure it is working properly and delivering a uniform spray pattern. As the volume of spray mixture decreases per acre, the importance of accurate calibration and uniform application increases. Avoid over-application, misapplication, and boom and spray swath overlapping that will increase spray dosage. (Crop injury may occur as a result.) Avoid spray skips and gaps which allow weeds to grow in untreated soil. Do not apply when weather conditions favor spray drift and/or when sensitive or cool season crops, such as cole crops, onions, peas, or strawberries are present in adjacent fields or in areas where wheat is growing in coarse textured soils.

APPLICATION: Uniform application is essential for satisfactory weed control. Accurate calibration of fertilizer application equipment is essential for uniform distribution to the soil surface. The correct method of application is to apply 1/2 the specified rate and overlap 50 percent or to double apply by splitting the middles to obtain the best distribution pattern. If fertilizer materials are excessively dusty, use diesel oil or other suitable additive to reduce dust prior to impregnation as dusty fertilizer will result in poor distribution during application. Crop injury and/or poor weed control may occur where the impregnated fertilizer is not uniformly applied.

INCORPORATION AND COMBINATION USES: When DIMETRIC DF 75% is to be used in combination with another herbicide, follow directions on the label for combinations, rates, crops, incorporation, and special precautions.

Refer in the label for tank mix information.

Limitations, Restrictions, and Exceptions

FOR WEED CONTROL IN A FALLOW ROTATION WITH BARLEY AND WHEAT

(Colorado, Kansas, Montana, Nebraska, and Wyoming Only)

DIMETRIC DF 75% may be applied to provide weed control during the fallow period after wheat or barley harvest or in the Spring before planting of Winter wheat or barley. Mechanical tillage or the application of a contact herbicide may be required to control weeds germinating prior to seeding of Winter wheat or barley.

For specific application information see the “General Information” section in the front of the label.

Where weed growth is present at application time, DIMETRIC DF 75% should be applied with Gramoxone, Roundup, or other contact herbicide. Refer to the other product label registered for additional directions, rates, and weed species controlled. Do not plant crops in treated areas earlier than 10 months following Fall applications. Do not rotate any crop not listed on the label for 18 months following application of DIMETRIC DF 75%.

AFTER HARVEST APPLICATION (Fall Fallow): DIMETRIC DF 75% may be applied to the stubble after harvest in the Fall. Apply 5/6 to 1 lb per acre broadcast before weeds emerge. Use the higher rate for longer weed control or for weeds designated as requiring the higher rate for control. Rainfall (1/2 inch or more) is necessary for herbicide activation.

Precautions and Restrictions: Do not graze treated fields.

Do not plant Spring seeded cereals following Fall applications for fallow. Where DIMETRIC DF 75% was applied in the Fall, do not apply DIMETRIC DF 75% in the Spring.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field_rates 0](#)

-

Restricted Entry Interval

12 hours

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

Timings

Post-harvest

Preemergence (Weed)