

WINTER WHEAT AND TRITICALE - FALL APPLICATION (WEEDS SUPPRESS)

General Information

Product Information

Use PowerFlex herbicide as a postemergence herbicide for the control of annual grass and broadleaf weeds in winter wheat and triticale. PowerFlex rapidly stops growth of susceptible weeds. However, typical symptoms (discoloration) of controlled or suppressed weeds may not be noticeable for 1 to 2 weeks after application, depending upon growing conditions and weed susceptibility. Degree of control and duration of effect are dependent upon weed sensitivity, weed size, crop competition, growing conditions at and following treatment, and spray coverage.

Use Precautions and Restrictions

When applying this product in tank mix combination, follow all applicable use directions, precautions, and limitations on each manufacturer's label.

Chemigation: Do not apply this product through any type of irrigation system.

Do not apply PowerFlex directly to, or otherwise permit it to come into direct contact with, susceptible crops or desirable plants including alfalfa, barley, canola, beans, cotton, flowers, grapes, lettuce, lentils, mustard, oats, peas, potatoes, radishes, soybeans, sugar beets, sunflowers, tobacco, tomatoes, vegetables, or other desirable broadleaf crops or ornamental plants. Do not permit spray mists containing PowerFlex to drift onto such plants.

Do not apply to crops underseeded with legumes.

Ground Applications: To minimize spray drift, apply PowerFlex in a total spray volume of 10 gallons or more per acre using spray equipment designed to produce large droplet, low pressure sprays. Refer to the spray equipment manufacturer's directions for detailed information on nozzle types, arrangement, spacing and operating height and pressure. Apply spot treatments only with a calibrated boom to prevent over application. Operate equipment at spray pressures no greater than is necessary to produce a uniform spray pattern. Operate the spray boom no higher

than is necessary to produce a uniformly overlapping pattern between spray nozzles. Do not apply with hollow cone-type insecticide nozzles or other nozzles that produce a fine-droplet spray.

Aerial Application: To minimize spray drift, apply PowerFlex in a total spray volume of 5 gallons or more per acre. Drift potential is lowest between wind speeds of 2 to 10 mph. However, many factors, including droplet size and equipment type, determine drift potential at any given speed. Avoid applications below 2 mph due to variable wind direction and high potential for temperature inversion. Minimize spray drift from aerial applications by applying a coarse spray at spray boom pressure no greater than 30 psi; by using straight-stream nozzles directed straight back; and by using a spray boom no longer than 3/4 of the rotor or wing span of the aircraft. Evaluate spray pattern and droplet size distribution by applying sprays containing a water-soluble dye marker or appropriate drift control agents over a paper tape (adding machine tape). Mechanical flagging devices may also be used.

Do not apply under conditions of a low level air temperature inversion. A temperature inversion is characterized by little or no wind and lower air temperature near the ground than at higher levels. The behavior of smoke generated by an aircraft-mounted device or continuous smoke column released at or near site of application will indicate the direction and velocity of air movement. A temperature inversion is indicated by layering of smoke at some level above the ground and little or no lateral movement.

Resistance Management

PowerFlex is an ALS mode of action (Group 2) herbicide. Any weed population may contain or develop plants naturally resistant to this product and other ALS herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. PowerFlex will not control known ALS (Group 2) resistant biotypes of labeled weeds. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance management strategies should be followed.

To delay herbicide resistance:

- For best resistance management stewardship, do not use more than once per season.
- Where possible, rotate the use of PowerFlex or other ALS herbicides with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted.
- Herbicide use should be based upon an IPM program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical), cultural, biological and other chemical control practices.
- Monitor treated weed populations for resistance development.
- Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment and planting clean seed.
- Contact your local extension specialist or certified crop advisers for any additional pesticide resistance management and/or integrated weed management requirements for specific crops and weed biotypes.

Application Directions

Application Timing

Apply PowerFlex postemergence to the main flush of actively growing weeds according to the target weed stage shown on the label. Extreme growing conditions such as drought, temperatures near or below freezing prior to, at, or following time of application may reduce weed control and increase the risk of crop injury at all stages of growth.

Warm, moist growing conditions promote active weed growth and enhance the activity of PowerFlex by allowing maximum foliar uptake and contact activity. Weeds hardened off by cold weather or drought stress may not be adequately controlled or suppressed and re-growth may occur. For best results, ensure thorough spray coverage of target weeds.

If foliage is wet at the time of application, control may be decreased. Applications of

PowerFlex are rainfast within 4 hours after application.

Refer to the Supplemental Label for the Crop Rotation Intervals for Use Only in the States of Idaho, Oregon, and Washington.

Limitations, Restrictions, and Exceptions

WINTER WHEAT AND TRITICALE

Apply 3.5 oz of PowerFlex per acre in either fall or spring to actively growing winter wheat and triticale from the 3-leaf to jointing stage (Zadoks scale 31) according to the application timings shown in the table entitled Weeds Controlled (C) or Suppressed (S) above. Treat after the majority of weeds have emerged. Best results are obtained when application is made to weeds that are actively growing.

Occasionally, slight yellowing or height reduction may be observed in the treated crop. These transient symptoms disappear within 14 days with no reduction to yield. Do not apply to crops suffering from drought, waterlogged soils, nutrient deficiency or exposed to frost or other agronomic factors affecting plant growth. Do not use on wheat or triticale varieties that are sensitive to ALS herbicides.

An independent liquid ammonium nitrogen fertilizer application made 7 days before or after an application of PowerFlex may result in transient leaf burn or stunting. Do not make a liquid fertilizer application during this period unless the risk of crop response is acceptable.

Crop Specific Use Restrictions:

- Preharvest Interval: Do not apply within 60 days of harvest.
- Do not graze the treated crop within 7 days following application.
- Do not cut the treated crop for hay within 28 days following application.
- Do not apply a product containing organophosphates for five days before or five days after an application of PowerFlex.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

60 days

Rates

[field_rates 0](#)

•

Restricted Entry Interval

12 hours

Timings

[Postemergence \(Crop\)](#)

[Postemergence \(Weed\)](#)

[fall](#)