

# **GRASSES GROWN FOR SEED - ANNUAL AND PERENNIAL WEEDS**

## General Information

### USE PRECAUTIONS

Unless otherwise noted under individual DIRECTIONS section, for aerial application, apply the recommended amount in a minimum of 2 gallons of water per acre. For ground application, apply the recommended amount in a minimum of 10 gallons of water per acre. Use more water for both methods when adverse growing conditions are present. DO NOT apply with high spray pressures, hollow cone or other nozzle types that produce small spray droplets which may drift. Avoid spray drift by making applications when conditions such as wind, air stability and temperature inversions are not a factor. The use of a suitable drift control agent at the proper rate will aid in the reduction of spray drift. Do not apply this product through any type of irrigation system.

### Limitations, Restrictions, and Exceptions

#### GRASSES GROWN FOR SEED

Do not use on alfalfa, clover, other legumes or newly seeded areas.

### DIRECTIONS

Apply to established grasses before the head comes into boot. Do not spray in boot stage. Apply to seedling grasses after grass has at least 5 leaves. Use the higher rate where weed stands are heavy.

Repeat treatment may be needed for less susceptible weeds. In some cases, Bent, Buffalo, Carpet and St. Augustine grasses may be injured by this treatment.

#### RESTRICTIONS AND LIMITATIONS FOR SEED GRASSES

Do not forage or graze treated areas for 7 days after application.

Do not apply more than 2 applications per year with a minimum retreatment interval of 21 days.

### Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field rates 0](#)

[field rates 1](#)

•

Restricted Entry Interval

12 hours

Timings

[To established grasses before the head comes into boot.](#)

[To seedling grasses after grass has at least 5 leaves.](#)