

# PECANS

## General Information

**AERIAL APPLICATION:** When applying SUPERIOR SPRAY OIL N.W. with aerial equipment, apply 2 to 5 gallons per acre. On young trees or oil susceptible varieties, use the lower rates. Use sufficient water and total gallonage per acre to give thorough coverage.

## USE PRECAUTIONS AND RESTRICTIONS

To avoid injury to trees, vines, fruit or foliage, observe the following precautions when using spray oils.

**WHEN USING SUPERIOR SPRAY OIL N.W. AS A TANK MIX WITH OTHER PRODUCTS, FOLLOW ALL LIMITATIONS, PRECAUTIONS AND RATE RECOMMENDATIONS WHICH APPEAR ON THE LABEL OF THE COMPANION PRODUCTS.**

- Do not apply this product through any type of irrigation system.
- Do not combine SUPERIOR SPRAY OIL N.W. with sulfur sprays.
- For Grapes: Do not apply sulfur within 10 days of an oil application. For all other crops, do not apply micronized sulfur within 10 days of an oil application and do not apply oil within 14 days of an application of wettable or dusting sulfur.
- After bloom, do not combine or follow an oil application with captan, carbaryl, dicofol, or dinocap as foliage or fruit injury may result.
- For both dormant and Summer oil sprays preceded or followed by applications of other materials that may not be compatible with oils, a sufficient interval needs to be maintained due to the potential for serious leaf and fruit bud injury leading to a reduction in fruit set.
- Do not use if emulsion breaks. If the emulsion should break in the spray tank, free oil will float to the surface causing excessive oil deposits at the end of each tank. Severe injury to trees may result if used under these conditions.
- When cool, slow drying conditions exist, labeled rates may increase the potential

for delayed bud development and/or leaf and fruit bud damage which may result in reduced fruit set. Use lower labeled rates or delay application until weather improves.

Do not apply when temperatures are likely to fall below freezing before the spray has a chance to dry.

Do not apply SUPERIOR SPRAY OIL N.W. when temperature is below 40° F or above 85° F.

**GROUND APPLICATION:** Apply using a nozzle height of no more than 4 feet above the ground or crop canopy.

Do not apply at wind speeds greater than 15 mph at the application site.

Apply as a medium or coarser spray (ASABE standard 572), and the minimum volume mean diameter (VMD) for spinning atomizer nozzles.

For airblast applications turn off outward pointing nozzles at row ends and when spraying outer row. To minimize spray loss over the top in orchard applications, spray must be directed into the canopy.

**AERIAL APPLICATION:** For aerial applications of agricultural products, do not release spray at a height greater than 10 feet above the ground, top of crops, or above the orchard canopy.

The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter.

When applications are made with a cross-wind, the swath will be displaced downwind. The applicator must compensate for this displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind. Leave at least one swath unsprayed at the downwind edge of the treated field.

Do not apply at wind speeds greater than 15 mph at the application site.

Apply as a medium or coarser spray (ASABE standard 572), and the minimum volume mean diameter (VMD) for spinning atomizer nozzles.

Limitations, Restrictions, and Exceptions

## PECANS

For control of obscure scale - Apply 3 to 3-1/2 gallons of SUPERIOR SPRAY OIL N.W. per 100 gallons of water. Application should be made during free dormancy and when temperature is between 50-70° F.

### Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

### Rates

[field\\_rates 0](#)

- 

Restricted Entry Interval

12 HOURS

### Timings

[During free dormancy and when temperature is between 50-70° F.](#)