

TANGERINE HYBRIDS - ENHANCE FRUIT SET AND YIELD

General Information

GENERAL INFORMATION

- Use only as directed. Read and thoroughly understand the label before making applications.
- The term 'grams of active ingredient per acre' is represented by 'g ai/acre' throughout this product label. See conversion table at the end of Directions for Use.
- Thoroughly spray all parts of the plant or crop to obtain the desired result.
- Prepare solution concentrations by mixing the required amount of product with water only in a clean empty spray tank.
- Dispose of any unused spray material at the end of the day. Refer to the Storage and Disposal section of the label for pesticide disposal instruction.
- When a range of rates is indicated, consult your local experimental station, distributor, or agricultural extension agent for the best program suited to your local conditions.
- Falgro 20SP works best when using water of neutral or slightly acidic pH. Ensure water pH is less than 8.5.
- Absorption of Falgro 20SP into the plant is greatest under slow drying conditions. Nighttime applications will be more effective when daytime conditions cause rapid drying. Re-apply Falgro 20SP if significant rain occurs within 2 hours of application.
- Falgro 20SP has a 0-day pre harvest interval (PHI).
- Do not apply using ULV application methods. For aerial applications spray volumes must be greater than 2 gallons per acre (10 gallons per acre for tree crops).
- Consult your local experimental station, distributor, or agricultural extension agent regarding the compatibility of gibberellic acid with other compounds.

Limitations, Restrictions, and Exceptions

SPRAYING GUIDE FOR CITRUS

Spray volume, quality and direction are chosen to ensure thorough wetting of all fruit. Do not exceed maximum application rates. Do not apply to trees of low vigor or suffering stress of any kind as this can cause severe fruit and/or leaf drop. Some drop of older leaves may occur after application of gibberellic acid. Never tank mix falgro 20SP with materials creating a high pH in the tank, e.g., white wash. The use of copper fungicides and/or oils within three weeks of a falgro 20SP application may result in significant leaf and/or fruit drop.

TANGERINE HYBRIDS

To enhance fruit set and yield: Application of falgro 20SP increases fruit set and yields in hybrids with pollination problems such as Minneola, Orlando, Robinson and Sunburst.

Make one application of 8 to 30 g ai/acre as a dilute spray during full bloom ensuring all foliage is wetted. A reduction in fruit size, retardation of color development and a slight increase in the drop of mature leaves may occur with this type of treatment.

Method

[Spray](#)

Rates

[field rates 0](#)

-

Restricted Entry Interval

4 hours

Timings

[During full bloom.](#)