

## **CHERRIES - RED TART (ALL APPROVED STATES EXCEPT CALIFORNIA) - 21 AND UP TREE AGE YEARS**

### General Information

#### GENERAL INFORMATION

- Use only as directed. Read and thoroughly understand the label before making applications.
- The term 'grams of active ingredient per acre' is represented by 'g ai/acre' throughout this product label. See conversion table at the end of Directions for Use.
- Thoroughly spray all parts of the plant or crop to obtain the desired result.
- Prepare solution concentrations by mixing the required amount of product with water only in a clean empty spray tank.
- Dispose of any unused spray material at the end of the day. Refer to the Storage and Disposal section of the label for pesticide disposal instruction.
- When a range of rates is indicated, consult your local experimental station, distributor, or agricultural extension agent for the best program suited to your local conditions.
- Falgro 20SP works best when using water of neutral or slightly acidic pH. Ensure water pH is less than 8.5.
- Absorption of Falgro 20SP into the plant is greatest under slow drying conditions. Nighttime applications will be more effective when daytime conditions cause rapid drying. Re-apply Falgro 20SP if significant rain occurs within 2 hours of application.
- Falgro 20SP has a 0-day pre harvest interval (PHI).
- Do not apply using ULV application methods. For aerial applications spray volumes must be greater than 2 gallons per acre (10 gallons per acre for tree crops).
- Consult your local experimental station, distributor, or agricultural extension agent regarding the compatibility of gibberellic acid with other compounds.

## Limitations, Restrictions, and Exceptions

### CHERRIES - RED TART (All approved states except California)

Application of Falgro 20SP maintains and extends high fruit bearing and reduces the occurrence of 'blind' nodes. The beneficial effect of Falgro 20SP application is not apparent until 2 - 3 years after initial application and is dependent on annually repeated applications. Make a single application 14 - 28 days after bloom when 3 - 5 terminal leaves are fully expanded or at least 1 - 3 inches of terminal growth have occurred. Apply Falgro 20SP according to the table below as a dilute spray in 100 - 150 gallons of water/acre. Lower water volumes can be used but it is extremely important not to overdose.

These application rates are based on tree vigor. Rates must be adjusted to the vigor expressed in each orchard. Use the higher rates for trees of low vigor and the lower rates for trees of high vigor. Applications cannot overcome the effects of nutritional, water, pest, disease or other stress on trees. Excessive application will increase vegetative growth at the expense of fruit production the following year.

#### Method

[Spray](#)

#### Rates

[field rates 0](#)

- 

#### Restricted Entry Interval

4 hours

#### Timings

[14 - 28 days after bloom when 3 - 5 terminal leaves are fully expanded or at least 1 - 3 inches of terminal growth have occurred.](#)