

## **VEGETABLE CROPS - CARROTS (FRESH AND PROCESSING)**

### General Information

#### GENERAL INFORMATION

- Use only as directed. Read and thoroughly understand the label before making applications.
- The term 'grams of active ingredient per acre' is represented by 'g ai/acre' throughout this product label. See conversion table at the end of Directions for Use.
- Thoroughly spray all parts of the plant or crop to obtain the desired result.
- Prepare solution concentrations by mixing the required amount of product with water only in a clean empty spray tank.
- Dispose of any unused spray material at the end of the day. Refer to the Storage and Disposal section of the label for pesticide disposal instruction.
- When a range of rates is indicated, consult your local experimental station, distributor, or agricultural extension agent for the best program suited to your local conditions.
- Falgro 20SP works best when using water of neutral or slightly acidic pH. Ensure water pH is less than 8.5.
- Absorption of Falgro 20SP into the plant is greatest under slow drying conditions. Nighttime applications will be more effective when daytime conditions cause rapid drying. Re-apply Falgro 20SP if significant rain occurs within 2 hours of application.
- Falgro 20SP has a 0-day pre harvest interval (PHI).
- Do not apply using ULV application methods. For aerial applications spray volumes must be greater than 2 gallons per acre (10 gallons per acre for tree crops).
- Consult your local experimental station, distributor, or agricultural extension agent regarding the compatibility of gibberellic acid with other compounds.

#### Limitations, Restrictions, and Exceptions

## SPRAYING GUIDE FOR VEGETABLE CROPS

### CARROTS (Fresh and Processing)

Apply falgro 20SP to delay leaf senescence and reduce the incidence of infection by *Alternaria dauci*. Make first application of 1 to 6 g ai/acre 4–6 weeks after emergence using commercial ground or aerial equipment with spray concentrations of 20-30 ppm. To achieve desired foliar recovery in severe disease situations or cool weather reapply in 14 days. Do not make more than two applications per crop cycle. Dilutions of greater concentrations can increase the risk of excessive top growth, particularly with a second application.

#### Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

#### Rates

[field\\_rates 0](#)

•

#### Restricted Entry Interval

4 hours

#### Timings

[4-6 weeks after emergence.](#)