

VEGETABLE CROPS - SPINACH - FALL AND OVERWINTERED (ALL APPROVED STATES EXCEPT CALIFORNIA)

General Information

GENERAL INFORMATION

- Use only as directed. Read and thoroughly understand the label before making applications.
- The term 'grams of active ingredient per acre' is represented by 'g ai/acre' throughout this product label. See conversion table at the end of Directions for Use.
- Thoroughly spray all parts of the plant or crop to obtain the desired result.
- Prepare solution concentrations by mixing the required amount of product with water only in a clean empty spray tank.
- Dispose of any unused spray material at the end of the day. Refer to the Storage and Disposal section of the label for pesticide disposal instruction.
- When a range of rates is indicated, consult your local experimental station, distributor, or agricultural extension agent for the best program suited to your local conditions.
- Falgro 20SP works best when using water of neutral or slightly acidic pH. Ensure water pH is less than 8.5.
- Absorption of Falgro 20SP into the plant is greatest under slow drying conditions. Nighttime applications will be more effective when daytime conditions cause rapid drying. Re-apply Falgro 20SP if significant rain occurs within 2 hours of application.
- Falgro 20SP has a 0-day pre harvest interval (PHI).
- Do not apply using ULV application methods. For aerial applications spray volumes must be greater than 2 gallons per acre (10 gallons per acre for tree crops).
- Consult your local experimental station, distributor, or agricultural extension agent regarding the compatibility of gibberellic acid with other compounds.

Limitations, Restrictions, and Exceptions

SPRAYING GUIDE FOR VEGETABLE CROPS

SPINACH – FALL AND OVERWINTERED (All approved states except California)

Application of falgro 20SP improves the quality, increases yield and aids harvest. Make one application of 6 to 10 g ai/acre 10–18 days before each anticipated harvest in 10–50 gallons of water if applied using ground equipment, or a minimum of 5–10 gallons of water/acre if sprayed by air. Ideally, spray in the early morning when dew is present on the crop and daytime temperatures are between 40 –70 F. Maximum benefit is seen from this treatment when low temperatures would limit the growth of untreated spinach.

Do not apply to spring sown spinach. Do not treat spinach after midwinter or if temperatures are expected to exceed 75 F within several days of application as this can induce bolting.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field rates 0](#)

-

Restricted Entry Interval

4 hours

Timings

[10–18 days before each anticipated harvest.](#)