

RICE SEED TREATMENT

General Information

GENERAL INFORMATION

- Use only as directed. Read and thoroughly understand the label before making applications.
- The term 'grams of active ingredient per acre' is represented by 'g ai/acre' throughout this product label. See conversion table at the end of Directions for Use.
- Thoroughly spray all parts of the plant or crop to obtain the desired result.
- Prepare solution concentrations by mixing the required amount of product with water only in a clean empty spray tank.
- Dispose of any unused spray material at the end of the day. Refer to the Storage and Disposal section of the label for pesticide disposal instruction.
- When a range of rates is indicated, consult your local experimental station, distributor, or agricultural extension agent for the best program suited to your local conditions.
- Falgro 20SP works best when using water of neutral or slightly acidic pH. Ensure water pH is less than 8.5.
- Absorption of Falgro 20SP into the plant is greatest under slow drying conditions. Nighttime applications will be more effective when daytime conditions cause rapid drying. Re-apply Falgro 20SP if significant rain occurs within 2 hours of application.
- Falgro 20SP has a 0-day pre harvest interval (PHI).
- Do not apply using ULV application methods. For aerial applications spray volumes must be greater than 2 gallons per acre (10 gallons per acre for tree crops).
- Consult your local experimental station, distributor, or agricultural extension agent regarding the compatibility of gibberellic acid with other compounds.

Limitations, Restrictions, and Exceptions

RICE SEED TREATMENT

Application of falgro 20SP as a seed treatment promotes germination and emergence of semi-dwarf and tall rice varieties but can only be applied to seed intended for drilling or dry broadcast. Do not apply to rice used in a 24-hour soak prior to broadcasting.

Falgro 20SP can be applied using mist-treatment equipment but best effect is achieved using the higher water volume. Half fill the seed treatment tank with water then add the required amount of falgro 20SP and mix thoroughly while adding any further seed treatment materials before making up to the desired final water volume. An EPA approved dye must be added to treated seed to prevent inadvertent use for food, feed or oil purposes. Falgro 20SP is compatible with most commonly used seed treatments, standard dyes and binding agents. However users should check to ensure adequate physical compatibility and mixing characteristics.

Method

[Seed Treatment](#)

Rates

[field rates 0](#)

-

Restricted Entry Interval

4 hours

Timings

[N.A.](#)