

## **BARLEY - APHID, ETC.**

### General Information

#### GENERAL INSTRUCTIONS

Use low rate under light to moderate infestation. Higher rates should be used under heavy insect pressure. The rate of application is variable according to insect pressure, timing of spray and field scouting. Do not exceed maximum allowable rate.

Under hot and dry conditions, increase the spray volume and add appropriate spray adjuvant, but only to those crops which have previously demonstrated a tolerance to the adjuvant(s) under these conditions.

#### Preventive Use

For cutworm, armyworm, or stalk borer control, Mustang insecticide may be applied before, during, or after planting. For soil-incorporated applications, use higher rates for improved control. Do not exceed maximum allowable rate.

Refer the specified label of Maximum Seasonal Usage and PHI for all Crops.

#### Limitations, Restrictions, and Exceptions

#### BARLEY

Apply as required by scouting. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds. Do not exceed maximum allowable rate.

Apply by ground or air equipment using sufficient water to obtain full coverage of foliage (minimum of 10 gallons by ground and 2 gallons by air).

For chinch bug control, begin applications when bugs migrate from small grains or grass weeds. Apply sufficient spray volume to penetrate the soil/stem interface, leaf collars, and sheaths.

Do not make applications less than 14 days apart.

Aphid spp, Whitefly spp, Greenbug: Aids in control. See resistance statement under

Directions For Use section.

Armyworm, Beet: See resistance statement under Directions For Use section.

Wheat Stem Sawfly (adult): Aids in control.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

14 days

Rates

[field\\_rates 0](#)

[field\\_rates 1](#)

•

Restricted Entry Interval

12 hours

Timings

[When insect populations reaching locally determined economic threshold levels.](#)