

## **CORN, FIELD - CORN EARWORM, ETC. (30 DAYS FOR GRAIN AND STOVER)**

### General Information

#### GENERAL INSTRUCTIONS

Use low rate under light to moderate infestation. Higher rates should be used under heavy insect pressure. The rate of application is variable according to insect pressure, timing of spray and field scouting. Do not exceed maximum allowable rate.

Under hot and dry conditions, increase the spray volume and add appropriate spray adjuvant, but only to those crops which have previously demonstrated a tolerance to the adjuvant(s) under these conditions.

#### Preventive Use

For cutworm, armyworm, or stalk borer control, Mustang insecticide may be applied before, during, or after planting. For soil-incorporated applications, use higher rates for improved control. Do not exceed maximum allowable rate.

Refer the specified label of Maximum Seasonal Usage and PHI for all Crops.

#### Limitations, Restrictions, and Exceptions

#### CORN, FIELD

Make applications when insect populations reach economic threshold levels. Refer to local Cooperative Extension Pest Management Guidelines and/or scouting results. Do not exceed maximum allowable rate.

Apply by air or by ground equipment using sufficient water to obtain full coverage of foliage (minimum of 2 gallons per acre by air and 10 gallons per acre by ground).

Direct spray to the base of plant. Repeat applications at 3 to 5 day intervals if needed. Mustang may only suppress heavy infestations and/or subsequent migrations.

Corn Earworm, Western Bean Cutworm: For control before the larva bores into the plant stalk or ear.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

30 days

Rates

[field\\_rates 0](#)

[field\\_rates 1](#)

•

Restricted Entry Interval

12 hours

Timings

[When insect populations reaching locally determined economic threshold levels.](#)