

# **SORGHUM GRAIN AND MILLET - APHID, ETC. (45 DAYS FOR FORAGE)**

## General Information

### GENERAL INSTRUCTIONS

Use low rate under light to moderate infestation. Higher rates should be used under heavy insect pressure. The rate of application is variable according to insect pressure, timing of spray and field scouting. Do not exceed maximum allowable rate.

Under hot and dry conditions, increase the spray volume and add appropriate spray adjuvant, but only to those crops which have previously demonstrated a tolerance to the adjuvant(s) under these conditions.

### Preventive Use

For cutworm, armyworm, or stalk borer control, Mustang insecticide may be applied before, during, or after planting. For soil-incorporated applications, use higher rates for improved control. Do not exceed maximum allowable rate.

Refer the specified label of Maximum Seasonal Usage and PHI for all Crops.

### Limitations, Restrictions, and Exceptions

### SORGHUM GRAIN

Apply as required by scouting. Timing and frequency of applications should be based upon insect populations reaching locally determined economic threshold levels. Do not exceed maximum allowable rate.

Apply by ground or air equipment using sufficient water to obtain full coverage of foliage (minimum of 10 gallons by ground and 2 gallons by air). The addition of one to two quarts of emulsified oil per acre to the spray solution may improve spray deposition and insect control.

For chinch bug control, begin applications when bugs migrate from small grains or grass weeds to small sorghum. Direct spray to the base of plants with sufficient spray volume to penetrate the soil/stem interface, leaf collars, and sheaths.

Do not make applications less than 10 days apart.

Aphid spp: Aphid control may be variable depending on species present and host-plant relationships. See resistance statement under \"Directions For Use\" section.

Armyworm, Beet: See resistance statement under \"Directions For Use\" section.

Lesser Cornstalk Borer: See resistance statement under \"Directions For Use\" section

Thrips spp.: See resistance statement under \"Directions For Use\" section. Aids in Control

#### Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

45 days

#### Rates

[field\\_rates 0](#)

[field\\_rates 1](#)

•

Restricted Entry Interval

12 hours

#### Timings

[chinch bug: when bugs migrate from small grains or grass weeds to small sorghum. When insect populations reaching locally determined economic threshold levels.](#)