

STONE FRUIT - AMERICAN PLUM BORER, ETC.

General Information

GENERAL INSTRUCTIONS

Use low rate under light to moderate infestation. Higher rates should be used under heavy insect pressure. The rate of application is variable according to insect pressure, timing of spray and field scouting. Do not exceed maximum allowable rate.

Under hot and dry conditions, increase the spray volume and add appropriate spray adjuvant, but only to those crops which have previously demonstrated a tolerance to the adjuvant(s) under these conditions.

Preventive Use

For cutworm, armyworm, or stalk borer control, Mustang insecticide may be applied before, during, or after planting. For soil-incorporated applications, use higher rates for improved control. Do not exceed maximum allowable rate.

Refer the specified label of Maximum Seasonal Usage and PHI for all Crops.

Limitations, Restrictions, and Exceptions

STONE FRUIT

Apply as required by scouting. Timing and frequency of applications should be based upon insect populations reaching locally determined economic threshold levels. Do not exceed maximum allowable rate.

Apply by ground or air equipment using sufficient water to obtain full coverage of foliage (for ground application use a minimum of 20 gallons for concentrate spray or a minimum of 100 gallons for dilute spray; for air application use a minimum of 10 gallons).

Do not make applications less than 7 days apart.

Avoid applications when honey bees are actively foraging by applying during the early morning or evening hours.

Do not apply as a ULV spray.

Do not feed or allow livestock to graze on cover crops from treated orchards.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

14 days

Rates

[field_rates 0](#)

[field_rates 1](#)

•

Restricted Entry Interval

12 hours

Timings

[When insect populations reaching locally determined economic threshold levels.](#)