

GRAPE

General Information

General Precautions and Restrictions: Do not allow adults, children, or pets to enter the treated area until sprays have dried. Do not apply this product in a way that will contact adults, children, or pets, either directly or through drift.

Limitations, Restrictions, and Exceptions

APPLICATION DIRECTIONS

Apply recommended amount using ground equipment, aircraft, or hand duster. Apply at the first signs of infestation and good leaf coverage is essential. Repeat at weekly intervals when needed to maintain control.

C-O-C-S 15 Sulfur 25 Dust is a product containing sulfur and copper oxychloride sulfate, a fixed neutral copper.

Application of material containing sulfur should not be made just prior to or during periods of excessively high temperatures.

GRAPES: Powdery Mildew, Bunch Rots—Use 19.5 pounds per acre. Apply at the time of berries begin to touch. For disease control the material must be applied before the disease appears.

Note: Grapes have a maximum per application rate of 19.5 lbs product/A; a maximum annual rate of 20.0 lbs product/A; and a minimum retreatment interval of 3 days.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field rates 0](#)

-

Restricted Entry Interval

48 hours

Timings

[At the time of berries begin to touch.](#)