

FOR CRACK AND CREVICE USE

General Information

Use Restrictions

- DO NOT reapply to treated area more than every four weeks.
- Use only in well-ventilated conditions.
- DO NOT contaminate food or food contact surfaces.
- In hospitals and nursing homes, remove patients before treating rooms. Allow treated areas to dry before returning patients to treated rooms.
- DO NOT use in schools while occupied. Allow treated areas to dry before allowing re-occupancy.
- DO NOT contact treated areas until spray has dried.
- Remove pets and cover or remove pet food/water and fish bowls prior to application.
- In zoos, remove animals and cover or remove animal food/water prior to application.
- DO NOT make general surface or space applications such as, for pantry pests in cupboards, cabinets, or in containers.
- Applications in food-handling areas of foodhandling establishments are limited to spot and/or crack and crevice treatments. The food-handling area may not be in operation during application. Exposed food must be covered or removed prior to application.
- DO NOT apply directly to food, food packaging, or food contact surfaces.
- After application, all food contact surfaces in the treated areas must be washed with an effective cleaning compound followed by a potable water rinse before resuming food operations.
- DO NOT make surface applications to mattresses or bedding. Mattress treatments

must be confined to the seams, folds and edges only. Remove linens and wash before reuse. Allow to dry before remaking bed.

Limitations, Restrictions, and Exceptions

For Crack and Crevice use: Press down the spray tip to release product while moving the spray can along length of treatment area. Apply at a rate of 2 sec/ln ft.

For indoor and exterior use as a crack and crevice or spot treatment only in the following areas to kill, on direct spray contact, the listed pests in the following areas: houses, apartments, meat, fish, poultry and egg processing and packaging plants, food/feed handling establishments (an area or place other than a private residence in which food is held, processed, prepared and/or served), and nonfood/feed handling areas of food handling establishments (garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage [after canning or bottling]), and commercial, institutional, warehousing establishments, and zoos.

Apply as a spot and/or crack and crevice and/or void treatment where pests harbor, nest, trail, run, and/or breed or hide. Treat into voids or to surfaces beneath cabinets, furniture, spaces between equipment, expansion joints, surfaces behind sinks, floor drains (to sewers), lockers, and water pipes.

Exterior treatments around the listed areas are limited to spot and/or crack and crevice treatments where pests enter around doors, windows, eaves, attic vents and holes in exterior walls where utilities enter.

To kill on direct spray contact drywood termites and subterranean termites in their above-ground galleries, workings or aerial nests within structures, locate existing openings or drill pilot holes to access termite tunnels, galleries, infested cavities or voids. Apply as a void treatment in holes to obtain thorough coverage of termites. Application of this product is not a substitute for a soil-applied termiticide. This product is not intended as a substitute for mechanical alteration, soil treatment or foundation treatment. DO NOT use to treat soil directly, When active infestations exist, get a professional inspection.

-