

GRASSES - ESTABLISHED LAWNS, GOLF COURSE, RESIDENTIAL TURF, SIMILAR TURF GRASSES AND SOD FARMS

General Information

PRODUCT INFORMATION

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

MCPA is effective on a large number of broadleaf weeds and is useful for controlling these weeds in certain crops. Several crops, such as flax, oats, and small grains underseeded to legumes, are more tolerant of MCPA than they are of 2,4-D. Crop varieties vary in response to MCPA, and some may be easily injured. Apply this product only to varieties known to be tolerant to MCPA. Injury to crops may occur from this pesticide. If you are not prepared to accept some degree of crop injury, do not use this product. If you are uncertain concerning tolerant varieties or local use situations that may affect crop tolerance to MCPA, contact your seed company or State Agricultural Extension Service for advice.

NOTE: Adding oil, wetting agent or other surfactant to the spray may reduce selectivity to crops, possibly resulting in crop injury.

RESTRICTIONS

See individual crop use directions for additional restrictions and maximum application rates. The minimum retreatment interval is 21 days.

Do not apply this product through any type of irrigation system.

Barley, oats, rye, wheat, flax and peas treated with MCPA may be replanted immediately with any crop specified on an MCPA label or any crop for which a residue tolerance exists for MCPA. For crops not listed on an MCPA label, or on crops for which no residue tolerances for MCPA have been established, a 60-day plantback interval must be observed.

This product will kill or seriously injure many desirable forms of vegetation. Do not

apply directly to flowers, fruits, grapes, tomatoes, ornamentals, cotton or other desirable plants. Vapors from this product may injure susceptible plants in the immediate vicinity. Do not apply when weather conditions favor drift from target area.

LIMITATIONS

Avoid use of small-diameter nozzles. Coarse sprays are less likely to drift. A spray thickening agent, such as Nalco-Trol, may be used with this product to aid in reducing spray drift. If used, follow all use recommendations and precautions on both product labels. Excessive amounts of this product in the soil may temporarily inhibit seed germination and plant growth. Determine air movement and direction before foliar application. Use a smoke generator or other means at or near the application site for the detection of air movement, air stability, or temperature inversions. Such a condition exists when there is little or no wind and air temperature is lower near the ground than at higher levels. Use appropriate drift control measures or avoid application when smoke is moving toward nearby desirable susceptible plants or sensitive areas.

WEEDS CONTROLLED

This product is effective on a large number of broadleaf weeds. On Mustard, Thistle, and White top or Hoary cress, it has superiority over 2,4-D.

SELECTIVE SPRAYING

NOTE: When using on flax, grain, grasslands, and pastures - do not forage or graze dairy and meat animals on treated areas within seven days of slaughter.

Limitations, Restrictions, and Exceptions

GRASSES

Established Lawns, Golf Courses, Residential Turf, Similar Turf Grasses and Sod Farms - Use 1 to 3.2 pints per acre in 10 to 120 gallons of water to give thorough coverage. Use higher rate for White top and Canada thistle. On smaller areas, use 1 fluid ounce (4 tablespoonfuls) mixed in 1 to 3 gallons of water and apply uniformly over 1,000 square feet. Do not exceed specified application dosages for any area. Avoid applying during excessively dry or hot periods unless irrigation (watering) is used before treatment. For best results, do not apply if rainfall is expected within 48 hours, and do not irrigate lawn for 48 hours. For optimum results, turf should not be mowed for 1 to 2 days before and after application. Reseed no sooner than 3 to 4

weeks after application of this product. Spring and fall are best times to treat. Do not use on lawns or creeping grasses, such as bent, except for spot spraying, nor on freshly-seeded turf until the grass has become well established, usually after the third mowing.

RESTRICTIONS AND LIMITATIONS FOR USE ON GOLF COURSES, RESIDENTIAL TURF, LAWNS, TURFGRASSES AND SOD FARMS

Do not apply more than 6.4 pints of this product (3.0 lb ae) per acre per year.

Do not apply more than 3.2 pints of this product (1.5 lbs ae) per acre per application.

Do not apply more than 2 applications per year with a minimum retreatment interval of 21 days.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field_rates 0](#)

•

Restricted Entry Interval

48 hours

Timings

[Postemergence \(Weed\)](#)