

CITRUS - WHITEFLIES AND CITRUS PSYLLID

General Information

DIRECTIONS FOR DILUTION UNLESS STATED

DILUTE APPLICATION: Ground Application For Field and Vegetable Crops: Apply specified rate in 20.0 to 60.0 gallons of water per acre.

CONCENTRATE APPLICATION: Ground Application: Apply specified rate in not less than 5.0 gallons of water per acre. Orchard Application:

Apply equivalent per acre rates in 20.0 to 100 gallons of water per acre. Special concentrate equipment is necessary for these uses.

AIR APPLICATION: Apply specified rate in 5.0 to 20.0 gallons of water per acre. Orchard rates use equivalent per acre rate in not less than 10.0 gallons of water per acre. Do not use air application on pecans.

Do not apply this product through any type of irrigation system.

Limitations, Restrictions, and Exceptions

CITRUS

Do not apply when trees or substantial numbers of weeds in the grove are in bloom. Do not use on seedlings. Do not apply within 15 days of harvest, except do not apply within 45 days of harvest for scale control rates over 1 1/2 pints (0.5 pound active ingredient) per 100 gallons of water. Make no more than 2 applications to mature fruit. Do not enter groves or nurseries within 48 hours of application. Do not graze livestock on cover crops grown in treated orchards. Thorough coverage is necessary. For aerial application, apply in a minimum of 5.0 gallons of water.

The REI is 10 days; however, the REI is increased to 14 days in outdoor areas where the average annual rainfall is less than 25 inches per year.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field_rates 0](#)

[field_rates 1](#)

•

Restricted Entry Interval

10 days

Timings

[N.A.](#)