

BERRY CROPS - SUPPRESSION OF LEAF AND BERRY DISEASE (AERIAL APPLICATION)

General Information

GENERAL APPLICATION INSTRUCTIONS

Apply this product by various application methods, including foliar spray (aerial and ground), soil drench, soil incorporation and bare root dip. For foliar sprays, apply this product with sufficient water volumes for adequate coverage of foliage, according to crop and growth stage.

Make applications prior to disease development in conjunction with good cultural management practices. Do not exceed the use rates or apply more frequently than the specified interval or plant injury can occur. Do not apply to plants that are dormant or heat or moisture stressed. To avoid undesirable copper phytotoxicity, do not make foliar applications to plants treated with copper-based compounds at less than 20 day intervals unless instructed to do so by your crop consultant. Allow foliage to dry completely after application. Do not apply when conditions favor wet tissue for prolonged periods (>4 hours).

See label for tank mix information.

Limitations, Restrictions, and Exceptions

BERRY CROPS

Application rate: Apply 1-1.5 quarts/acre (2-3.5 L/ha) in a minimum of 20 gallons/acre (190 L/ha) of water.

Not registered for use in CA unless accompanied by a supplemental label.

Low Disease Pressure: Apply lower rate at the first onset of the disease. Repeat applications at 1-3 week intervals. Do not apply more than 6 times per crop cycle.

High Disease Pressure: Apply higher rate at the first onset of the disease. Repeat applications at 7- 10 day intervals. Do not apply more than 6 times per crop cycle.

Ideally Fungi-Phite is best used in combination with conventional registered

fungicides to increase the performance of the disease control program.

Method

[Broadcast/Foliar Air](#)

Rates

[field rates 0](#)

[field rates 1](#)

-

Restricted Entry Interval

4 hours

Timings

[At the first onset of the disease.](#)