

PEANUT - WIREWORM (SUPPRESSION)

General Information

Product Information

Govern 4E insecticide is an emulsifiable concentrate for use in listed crops. This product resists washoff once it is dry. Target pests and application rates are provided in the accompanying tables.

Use Precautions and Restrictions

Insect control may be reduced at low spray volumes under high temperature and wind conditions.

Some reduction in insect control may occur under unusually cool Conditions

Do not aeri ally apply this product in Mississippi.

Insecticide Resistance Management (IRM) Govern 4E contains a Group 1B insecticide. Insect/mite biotypes with acquired resistance to Group 1B may eventually dominate the insect/mite population if Group 1B insecticides are used repeatedly in the same field or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by Govern 4E or other Group 1B insecticides.

To delay development of insecticide resistance, the following practices are recommended:

- Avoid consecutive use of insecticides with the same mode of action (same insecticide group) on the same insect species.
- Use tank mixtures or premix products containing insecticides with different modes of action (different insecticide groups) provided the products are registered for the intended use.
- Base insecticide use on comprehensive Integrated Pest Management (IPM) programs.

- Monitor treated insect populations in the field for loss of effectiveness.
- Contact your local extension specialist, certified crop advisor, and or manufacturer for insecticide resistance management and/or IPM recommendations for the specific site and resistant pest problems.
- For further information or to report suspected resistance, call 800-258-3033.

Limitations, Restrictions, and Exceptions

Peanut

Apply to the soil surface as a preplant broadcast spray followed by immediate soil incorporation to a depth of 3 to 4 inches using a disc, field cultivator, or equivalent equipment. Use a minimum of 10 gpa total spray.

Specific Use Restrictions:

- The combined total of preplant and postplant applications of Govern 4E, Lorsban 15G or other product containing chlorpyrifos must not exceed 4 lb ai chlorpyrifos per acre per season.
- Maximum single application rate is 2 lb ai chlorpyrifos (4 pints of Govern 4E) per acre.
- Do not feed treated peanut forage or hay to meat or dairy animals.
- Do not aerially apply this product in Mississippi.

Method

[Surface](#)

[Broadcast Spray](#)

Pre-Harvest Interval

21 days

Rates

[field rates 0](#)

•

Restricted Entry Interval

24 hours

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

Timings

[Preplant](#)