

WHEAT (FORAGE AND HAY) - ARMY CUTWORMS, ETC.

General Information

Product Information

Govern 4E insecticide is an emulsifiable concentrate for use in listed crops. This product resists washoff once it is dry. Target pests and application rates are provided in the accompanying tables.

Use Precautions and Restrictions

Insect control may be reduced at low spray volumes under high temperature and wind conditions.

Some reduction in insect control may occur under unusually cool Conditions

Do not aerially apply this product in Mississippi.

Insecticide Resistance Management (IRM) Govern 4E contains a Group 1B insecticide. Insect/mite biotypes with acquired resistance to Group 1B may eventually dominate the insect/mite population if Group 1B insecticides are used repeatedly in the same field or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by Govern 4E or other Group 1B insecticides.

To delay development of insecticide resistance, the following practices are recommended:

- Avoid consecutive use of insecticides with the same mode of action (same insecticide group) on the same insect species.
- Use tank mixtures or premix products containing insecticides with different modes of action (different insecticide groups) provided the products are registered for the intended use.
- Base insecticide use on comprehensive Integrated Pest Management (IPM) programs.

- Monitor treated insect populations in the field for loss of effectiveness.
- Contact your local extension specialist, certified crop advisor, and or manufacturer for insecticide resistance management and/or IPM recommendations for the specific site and resistant pest problems.
- For further information or to report suspected resistance, call 800-258-3033.

Limitations, Restrictions, and Exceptions

Wheat

Foliar Application

Apply using aerial (fixed wing or helicopter) or power-operated ground spray equipment. Mix the required dosage with water and apply in a minimum of 2 to 5 gpa finished spray volume for aerial equipment or 15 gpa for ground spray equipment. Apply when field counts indicate damaging pest populations are developing or present.

Chemigation: Govern 4E may be applied through sprinkler irrigation systems at specified broadcast application rates to control listed foliar pests. See Chemigation Application section for application instructions.

cutworms (suppression))

Pest-Specific Use Directions:

Control may be reduced under high temperature conditions (greater than 80°F), under dry soil conditions, or if larvae are more than 1/2 inch long.

Suppression should be expected under conditions of heavy pest populations or large worms.

Target application when eggs are near hatching and larvae is emerging as monitored by plant inspection.

For control of wheat midge, treat when 75% of the wheat heads have emerged from the boot and when midge adults are found in the crop (1 midge per 4 to 5 heads). If possible, apply in the late afternoon or early evening when temperatures exceed 50°F and wind speed is less than 7 mph.

Specific Use Restrictions:

- Maximum single application rate is 0.5 lb ai chlorpyrifos (1 pint of Govern 4E) per acre.
- Do not allow meat or dairy animals to graze or otherwise feed on treated forage within 14 days of application.
- Do not feed straw from treated wheat within 28 days of application.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

14 days

Rates

[field rates 0](#)

-

Restricted Entry Interval

24 hours

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

Timings

[When field counts indicate damaging pest populations are developing or present.](#)