

OUTDOOR / PERIMETER PESTS - FIRE ANTS

General Information

PRODUCT FACTS

- Two modes of action
- Contact activity & systemic action
- Quick knockdown & residual control
- Low odor
- Non-staining
- No phytotoxicity on a wide range of ornamentals

Temprid SC Insecticide

- Temprid SC Insecticide is a suspension concentrate formulation of two insecticide active ingredients, imidacloprid and b-cyfluthrin. One gallon of the product contains 2 lbs of imidacloprid and 1 lb of bcyfluthrin.
- Shake well before using.

Controls

- Indoor or General Household Pests
- Outdoor/Perimeter Pests
- Wood-Infesting Pests

Where To Apply

Apply where pests have been seen or found, or can find shelter.

- In and around buildings and structures
- On established landscape ornamentals around residential and commercial

properties

Dilution Rates

- When 0.27 fl oz (8 milliliters) is diluted in one gal of water, the combined active ingredient concentration (0.075%) contains 0.05% imidacloprid and 0.025% b-cyfluthrin.
- When 0.54 fl oz (16 milliliters) is diluted in one gal of water, the combined active ingredient concentration (0.15%) contains 0.10% imidacloprid and 0.05% b-cyfluthrin.

Mixing Information

- Mix the appropriate amount of Temprid SC Insecticide with water.
- Add product when filling spray tank with water; shake or agitate mixture.
- Diluted spray mixture can be stored overnight. Agitate before using.

How To Apply

- General surface (outdoors only), spot treatment, and void injections by mist, foam or crack & crevice sprays.
- Foliar application to established landscape ornamentals.

Re-Apply

- Every 7 to 10 days, if needed

Re-Entry

- Avoid contact with treated surfaces until dry.
- People and pets may re-enter after the treated area is dry.

Limitations, Restrictions, and Exceptions

OUTDOOR / PERIMETER PESTS

WHERE TO APPLY:

- Apply to surfaces on buildings, porches, patios and other structures, around doors and windows, eaves and attic vents, utility entry points, soffit areas and other exterior openings (including foundation cracks or drilled holes) where these pests enter the structure or where they have been seen or found, or can find shelter.
- To prevent infestation, treat the outside of the structure, to include walls (to a height of 2 to 3 feet), around doors, windows, and soffits where pests are active or may enter or hide. Treat the soil, turf or other ground covering adjacent to the structure (in a band up to 10 feet wide) where pests are seen, have been found, or can find shelter.
- In ant management programs, apply to flower, shrub or ornamental plant beds where ants may find food or forage.

USE RATE:

- 0.15%: 0.54 fl oz (16 mL) per gallon of water.

Always use 0.15% rate for fire ants, hoents, wasps and yellowjackets.

HOW TO APPLY:

- High volume applications using power sprayers: In power sprayers, adjust the rate of product dilution to apply 8 to 16 milliliters of Temprid SC Insecticide per 1,000 square feet in sufficient water to provide adequate coverage. For example, mixing 4.5 fluid ounces (133 milliliters) of the procdut in a 50 gallon tank, when applied in a volume of 3.0 gallons per 1,000 square feet, will apply 0.27 fluid ounces of the procdut per 1,000 square feet. (See label for the table).
- Apply enough dilution to adequately cover the area without excessive dripping or run-off. Volume can vary depending on the surface type treated. For example, mulch requires more spray than bare soil. Do not contact treated surfaces until dry.

WHEN TO APPLY:

- When pests are seen or found.
- Re-apply every 7 to 10 days if needed.
- Do not apply more than 32.0 fluid ounces of Temprid SC Insecticide per acre, per

year for landscape, lawn and ornamental applications.

- At the 8 milliliter per gallon use rate, this is a limit of 118 dilute gallons per acre, per year and at the 16 milliliter per gallon use rate, this is a limit of 59 dilute gallons per acre, per year for landscape, lawn, and ornamental applications.

APPLICATION RESTRICTIONS:

- Do not apply directly to impervious horizontal surfaces such as sidewalks, driveways, and patios except as a spot or crack and crevice treatment. During application, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. All outdoor applications, if permitted elsewhere on the label, must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses:

1. Applications to soil or vegetation, as listed on the label, around structures;
2. Applications to lawns, and other vegetation, as listed on the label;
3. Applications to the side of a building, up to a maximum height of 3 feet above grade;
4. Applications to underside of eaves, soffits, doors, or windows permanently protected from rainfall by a covering, overhang, awning, or other structure;
5. Applications around potential pest entry points into buildings, when limited to a surface band not to exceed one inch in width;
6. Applications made through the use of a coarse, low pressure spray to only those portions of surfaces that are directly above bare soil, lawn, turf, mulch or other vegetation, as listed on the label, and not over an impervious surface, drainage or other condition that could result in runoff into storm drains, drainage ditches, gutters, or surface waters, in order to control occasional invaders or aggregating pests.

SPECIFIC INSTRUCTIONS FOR PESTS (Fire Ants)

- Thoroughly drench or inject the entire mound. When using injection treatment, you must treat the bottom of the mound. Apply dilution or foam using a downward and upward motion of the injector tool. For complete distribution, inject in several points

throughout the mound.

Treat new mounds as they appear.

- Do not use in pastures or areas of food crop production.

Method

[Directed](#)

[Spray](#)

Rates

[field_rates 0](#)

[field_rates 1](#)

[field_rates 2](#)

•

Timings

[When pests are seen or found.](#)