

TURFGRASS - COOL SEASON GRASSES - BARNYARD GRASS, CHICKWEED, ETC.

General Information

GENERAL INFORMATION

This product is a selective herbicide that provides preemergence control of crabgrass and other susceptible annual grasses and broadleaf weeds in established lawns and ornamental turf. This product will not harm nearby established ornamentals when used according to label directions. In New York State this product may only be used by professional applicators at no more than 185 pounds (0.5 pounds active ingredient) per year and is prohibited from use in Nassau and Suffolk Counties.

PRECAUTIONS

Use of this product on golf greens is not recommended. DO NOT apply this product to Colonial bentgrass. For best results, cultural practices that disturb the soil, such as core, spike, hydro-aerification, or verticutting, and top dressing should be done before application of this product. This product should not be applied until turfgrasses have recovered from stresses related to these or other mechanical operations. Irrigation or rainfall within 7 days after application will improve the performance of this product.

Reseeding or Sprigging: Reseeding, overseeding, or sprigging of treated areas within 4 months after application of this product may inhibit the establishment of desirable turfgrass. When reseeding or overseeding, proper cultural practices such as soil cultivation, irrigation and fertilization should be followed. For best results, use mechanical or power seeding equipment (slit seeders) designed to give good seed-to-soil contact. This product may be used on seeded, sodded, or sprigged turfgrass that are well-established. Allow the turf to develop a good root system and uniform stand before application.

Use of this product on turfgrass not well-established, weakened by lack of moisture, chemical injury, insects, disease, or temperature extremes that cause stress will increase the potential for injury. Allow turfgrass weakened by these stresses to

recover before application of this product. DO NOT use clippings from treated turf for mulching around vegetables or fruit trees. DO NOT allow domestic animals to feed or forage on the treated turf.

This product may be applied as a single application or as a split application in the spring, summer or fall. DO NOT apply more than 185 pounds of this product per acre per application and no more than 555 pounds (185 pounds in New York) of this product per acre per year. Do not apply in Nassau and Suffolk Counties of New York.

Limitations, Restrictions, and Exceptions

TURFGRASS

- Use of this product on certain varieties of creeping bentgrass, such as "Carmen", "Cohansey", or "Seaside" will result in undesirable turf injury.

Not all varieties of creeping bentgrass have been tested.

- The following fine fescue varieties have been found to be sensitive to this product: "Atlanta", "Banner", "Beauty", "Bligart", "CF-2", "Enjoy", "HF-93", "Highlight", "Ivalo", "Jamestown", "Koket", "Majenta", "Mary", "Pennlawn", "Tamara", "Tatjana", "Waldorf", and "Waldina"

APPLICATION EQUIPMENT AND INSTRUCTIONS

- Apply this product with drop or rotary-type spreaders designed to apply granular herbicides. Calibrate application equipment prior to usage, and apply this product uniformly over the turfgrass area. A more uniform application can be made by spreading half of the required amount over the area and then applying the remaining half at a right angle to the previous, (see spreader settings for half rate). Avoid the use of spreaders that would apply this product in narrow rows or concentrated bands. Avoid streaking, skips, or overlaps during application. Check equipment frequently to ensure equipment is functioning properly and applying uniform distribution of granules.

APPLICATION DIRECTIONS

- This product provides preemergence control of listed annual grasses and broadleaf weeds when applied prior to their germination in established

turfgrass.

CONTROL OF ANNUAL GRASSES AND/OR ANNUAL BROADLEAF WEEDS

- This product should be applied prior to the germination of annual grasses and/or broadleaf weeds. Use the higher recommended rates where weed infestations are dense and/or applications are made early and longer residual control is desirable. For preemergence control and/or suppression of annual bluegrass and listed winter annual broadleaf weeds, apply this product prior to their germination and emergence. Use the higher recommended use rates where weed infestations are dense.

NOTE: The higher use rates are recommended where weed infestations are heavy and/or dense.

Method

[Broadcast](#)

Rates

[field_rates 0](#)

[field_rates 1](#)

•

Restricted Entry Interval

12 hours

Timings

[Preemergence \(Weed\)](#)