

OUTDOOR ORNAMENTALS - AZALEA SCALE, BAGWORM, ETC.

General Information

PRECAUTIONS AND RESTRICTIONS

BUFFER ZONES FOR AERIAL APPLICATION: When making a Non-ULV application with aerial application equipment, a minimum buffer zone of 25 feet must be maintained along any water body.

APPLICATIONS

Use higher rate when foliage is heavy or infestation is severe. Apply when pests first appear. Apply the following specified rates in sufficient water to thoroughly cover one acre. By ground, apply using a minimum of 10 gallons of water/A and by air apply using a minimum of 2 gallons of water/A (standard is 100 gallons of water for thorough coverage sprays). Do not apply orchard rates in less than 10 gallons of water per acre.

Limitations, Restrictions, and Exceptions

OUTDOOR ORNAMENTALS

Precaution: Before treating a large number of ornamental plants with FYFANON® 8 LB EMULSION alone or as a tank mixture with any other material, make a test application on a few plants and observe 7-10 days prior to treating large areas to reduce the possibility of plant injury.

COMMENTS:

- CAUTION: Avoid use on certain ferns including Boston, Maidenhair and Pteris, as well as some species of Crassula and Canaetri Juniper. 10-day minimum retreatment interval.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

field_rates 0

-

Restricted Entry Interval

12 hours

Timings

For Juniper, Fletch, Oak Kermes: When scale crawlers have settled on foliage.