

IMPATIENS (0.25 LB / 4350 SQ FT OR 0.1 / ACRE)

General Information

GENERAL INFORMATION

MilStop is a broad spectrum, contact foliar fungicide that controls a variety of diseases on terrestrial and indoor woody and herbaceous ornamental plants, and on greenhouse, nursery and agricultural crops. MilStop is applied using a sufficient volume of water to insure complete coverage of all stems and foliage.

NOTICE TO USER: Due to variability within species and in application techniques and equipment, and possible tank mix combinations, neither the manufacturer nor the seller has determined whether MilStop can be safely used on all varieties and species of ornamentals, greenhouse, and agricultural plants under all conditions. Therefore, the professional user should determine if MilStop can be used safely before broad usage by applying the prescribed use rate under the conditions expected to be encountered to a small test area. Any adverse effects should be visible within 14 days.

For curative disease control on sensitive crops apply at a rate of 2.5 pounds of MilStop per 100 gallons of water per week and increase the rate each week in 0.25 pound increments. Do not exceed a use rate of 5.0 pounds per acre per application.

SPRAY APPLICATION METHOD

Do not apply this product through any type of irrigation system. Do not apply directly to soil as a directed spray or drench. For best results, use a properly calibrated sprayer or similar equipment, and check equipment calibration frequently.

APPLICATION TIMING

Start application at first sign of disease. For best protection, repeat at one to two week intervals until conditions are no longer favorable for disease development. Shorten the interval during rainy weather or during periods of high relative humidity. For appropriate crops, apply MilStop as a post-harvest treatment to remaining foliage to decrease overwintering inoculum.

Limitations, Restrictions, and Exceptions

IMPATIENS - (0.25 LB / 4350 SQ FT OR 0.1 / ACRE)

During vegetative growth; discontinue spraying when blossom buds are visible.

Method

[N. A.](#)

Rates

[field_rates 0](#)

[field_rates 1](#)

•

Restricted Entry Interval

1 hour

Timings

[During vegetative growth.](#)