

BERRY CROPS

General Information

GENERAL INFORMATION

NUE Copper 5% is manufactured specifically as a liquid foliar fertilizer to be used to help prevent and correct copper deficiencies that may limit crop growth and yields. NUE Copper 5% is a supplement to an adequate and balanced soil fertility program.

NUE Copper 5% is a non-sulfate source of copper, allowing for blending with calcium products, specifically NUE Cal-8.

It is important to monitor crop needs and responses by maintaining an adequate tissue-testing program. When unfamiliar with the effects of foliar applications of minerals it is recommended to perform small-scale field trials.

DIRECTIONS FOR USE

Dilution and Spray Volumes - Always use enough water for uniform application and to prevent possible plant tissue damage to the intended crop from concentrated droplets. It is recommended to use a dilution of at least 20 parts of water to 1 part of NUE Copper 5%.

SUGGESTED APPLICATIONS

Optimum rate of application will vary between fields depending on soil and/or climatic conditions, stage of crop growth and size of plants.

Application rates are best determined by soil and plant analysis recommendations.

NUE Copper 5% can be used on many crops where foliar copper applications are commonly used. Consult your Bio-Gro dealer for specific recommendations.

TIMING - Application timing is likely the most important factor in determining the response and value of foliarly applied nutrients. Understanding the basic physiological functions of a crop is essential in planning proper nutrient timing.

Limitations, Restrictions, and Exceptions

BERRY CROPS

Applications may be repeated in 7 to 14 day intervals.

Method

[Foliar spray](#)

Rates

[field rates 0](#)

-

Timings

[After active vegetative growth begins.](#)