

FOLIAR APPLICATION: AERIAL APPLICATION

General Information

General Information

BRANDT LIGNO 7% Zn is recommended to prevent and correct zinc deficiencies of field, row, vegetable, fruit, tree, and vine crops. Applied alone, with pesticides, or with other fertilizers, this product may be applied directly to the soil or to the foliage of growing crops via ground sprayers, aircraft or fertigation systems.

Mixing and Handling Instructions

Shake, stir, or swirl contents before using. Put 1/3 to 2/3 of total required water volume in tank. Add recommended amount of this product before adding pesticides, adjuvants, or other fertilizers. Agitate until thoroughly mixed, then fill tank with remainder of water.

BRANDT LIGNO 7% Zn is compatible with most pesticides and fertilizers. However, if the desired combination has not been previously used, check for chemical mixture compatibility using a jar test prior to application.

Limitations, Restrictions, and Exceptions

FOLIAR APPLICATION: Actively growing plants with young leaves absorb foliarly applied BRANDT ORGANICS 7% Zn most efficiently. Immediate post-irrigation or post-rainfall application will enhance foliar uptake. Thorough coverage and wetting are needed for optimum foliar spray results.

Applications should be separated by a minimum of 7-14 days. AVOID foliar applications of this product when plants are under moisture stress. AVOID foliar applications to deciduous tree crops during bloom.

Aerial Applications: Up to 1 quart per 5 gallons of spray may be used, but more dilute solutions will improve coverage and results.

Method

[Broadcast/Foliar Air](#)

Rates

field_rates 0

-

Timings

Actively growing plants with young leaves .