

GRASS FORAGE, FODDER, AND HAY GROUP AND GRASS GROWN FOR SEED AND PASTURE AND RANGELAND - ALFALFA CATERPILLAR, ETC.

General Information

GENERAL INSTRUCTIONS

Use low rate under light to moderate infestation. Higher rates should be used under heavy insect pressure. The rate of application is variable according to insect pressure, timing of spray and field scouting. Do not exceed maximum allowable rate.

Under hot and dry conditions, increase the spray volume and add appropriate spray adjuvant, but only to those crops which have previously demonstrated a tolerance to the adjuvant(s) under these conditions.

Preventive Use

For cutworm, armyworm, or stalk borer control, Mustang insecticide may be applied before, during, or after planting. For soil-incorporated applications, use higher rates for improved control. Do not exceed maximum allowable rate.

Refer the specified label of Maximum Seasonal Usage and PHI for all Crops.

Limitations, Restrictions, and Exceptions

GRASS

Apply as insects appear in sufficient volume of water to ensure thorough coverage of foliage.

Use higher recommended dosage for increased pest pressure or for increased residual pest control. Do not exceed maximum allowable rate.

Apply in a minimum of 2 gallons of finished spray per acre by aerial equipment or 10 gallons per acre by ground equipment.

ULV oil spray application is prohibited. Higher volumes of finished spray may improve insect control under high temperatures, when foliage is dense and/or when

insect pressure is high.

Do not make applications less than 7 days apart for forage and hay; not less than 17 days for straw and seed screenings.

Do not spray livestock. Allow application to dry before letting livestock graze on treated area.

A maximum of 4.3 ounces of product or 0.05 pounds of active ingredient per acre may be applied per cutting.

For hay, a maximum of 17.2 ounces of product or 0.20 pounds of active ingredient per acre per season may be applied.

For forage, straw, and seed screenings, a maximum of 21.5 ounces of product or 0.25 pounds of active ingredient per acre per season may be applied.

Applications may be made up to 0 days for forage and hay; 7 days for straw and seed screenings.

Aphid control may be variable depending on species present and host-plant relationships.

Blue Alfalfa Aphid, Green Peach Aphid, Pea Aphid, Spotted Alfalfa Aphid: Aphid control may be variable depending on species present and host-plant relationships.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field_rates 0](#)

[field_rates 1](#)

•

Restricted Entry Interval

12 hours

Timings

[When insects appear.](#)