

GRASS FORAGE, FODDER, AND HAY GROUP AND GRASS GROWN FOR SEED AND PASTURE AND RANGELAND - ARMYWORM, ETC.

General Information

GENERAL INSTRUCTIONS

Use low rate under light to moderate infestation. Higher rates should be used under heavy insect pressure. The rate of application is variable according to insect pressure, timing of spray and field scouting. Do not exceed maximum allowable rate.

Under hot and dry conditions, increase the spray volume and add appropriate spray adjuvant, but only to those crops which have previously demonstrated a tolerance to the adjuvant(s) under these conditions.

Preventive Use

For cutworm, armyworm, or stalk borer control, Mustang insecticide may be applied before, during, or after planting. For soil-incorporated applications, use higher rates for improved control. Do not exceed maximum allowable rate.

Refer the specified label of Maximum Seasonal Usage and PHI for all Crops.

Limitations, Restrictions, and Exceptions

GRASS

Apply as insects appear in sufficient volume of water to ensure thorough coverage of foliage.

Use higher recommended dosage for increased pest pressure or for increased residual pest control. Do not exceed maximum allowable rate.

Apply in a minimum of 2 gallons of finished spray per acre by aerial equipment or 10 gallons per acre by ground equipment.

ULV oil spray application is prohibited. Higher volumes of finished spray may improve insect control under high temperatures, when foliage is dense and/or when insect pressure is high.

Do not spray livestock. Allow application to dry before letting livestock graze on treated area.

For hay, a maximum of 17.2 ounces of product or 0.20 pounds of active ingredient per acre per season may be applied.

For forage, straw, and seed screenings, a maximum of 21.5 ounces of product or 0.25 pounds of active ingredient per acre per season may be applied.

Applications may be made up to 0 days for forage and hay; 7 days for straw and seed screenings.

Aphid control may be variable depending on species present and host-plant relationships.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field rates 0](#)

[field rates 1](#)

•

Restricted Entry Interval

12 hours

Timings

[When insects appear.](#)