

SUNFLOWER, ETC. - FOLIAR USE - SUNFLOWER BEETLE, ETC.

General Information

GENERAL INSTRUCTIONS

Use low rate under light to moderate infestation. Higher rates should be used under heavy insect pressure. The rate of application is variable according to insect pressure, timing of spray and field scouting. Do not exceed maximum allowable rate.

Under hot and dry conditions, increase the spray volume and add appropriate spray adjuvant, but only to those crops which have previously demonstrated a tolerance to the adjuvant(s) under these conditions.

Preventive Use

For cutworm, armyworm, or stalk borer control, Mustang insecticide may be applied before, during, or after planting. For soil-incorporated applications, use higher rates for improved control. Do not exceed maximum allowable rate.

Refer the specified label of Maximum Seasonal Usage and PHI for all Crops.

Limitations, Restrictions, and Exceptions

SUNFLOWER

Apply with ground or air equipment using sufficient water and application methods to insure thorough coverage of foliage. Apply in a minimum of 2 gallons of finished spray per acre by aerial equipment or 10 gallons per acre by ground equipment. Begin applications when pest appears and repeat as necessary to maintain control. Do not make applications less than 7 days apart.

Use higher recommended dosage for increased residual pest control. Do not exceed maximum allowable rate.

Grazing: Do not graze livestock in treated areas or cut treated crops for feed

Avoid applications when honey bees are actively foraging by applying during the early morning or evening hours.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

30 days

Rates

[field_rates 0](#)

[field_rates 1](#)

•

Restricted Entry Interval

12 hours

Timings

[When pest appears.](#)