

BROCCOLI, BRUSSELS SPROUT, CABBAGE, AND CAULIFLOWER TO BE EXPORTED TO CANADA - ARIZONA AND CALIFORNIA

General Information

GENERAL INSTRUCTIONS

Use low rate under light to moderate infestation. Higher rates should be used under heavy insect pressure. The rate of application is variable according to insect pressure, timing of spray and field scouting. Do not exceed maximum allowable rate.

Under hot and dry conditions, increase the spray volume and add appropriate spray adjuvant, but only to those crops which have previously demonstrated a tolerance to the adjuvant(s) under these conditions.

Preventive Use

For cutworm, armyworm, or stalk borer control, Mustang insecticide may be applied before, during, or after planting. For soil-incorporated applications, use higher rates for improved control. Do not exceed maximum allowable rate.

Refer the specified label of Maximum Seasonal Usage and PHI for all Crops.

Limitations, Restrictions, and Exceptions

BROCCOLI

This use pattern is designed to meet the Canadian Default MRL. Follow all instructions, recommendations and restrictions on the Mustang Insecticide label with the following modifications for produce to be exported to Canada.

- Application Interval: Do not make applications less than 14 days apart.
- Do not use Mustang Max or Mustang Maxx or products containing cypermethrin or alpha-cypermethrin on crops treated with Mustang Insecticide (combined residues may exceed the Canadian Default MRL).

Method

[Broadcast/Foliar Air](#)

Broadcast/Foliar Ground

Pre-Harvest Interval

14 days

Rates

[field rates 0](#)

[field rates 1](#)

-

Restricted Entry Interval

12 hours

Timings

[N.A.](#)