

ORNAMENTAL PLANTS - BROAD MITE, EUROPEAN RED MITE, ETC.

General Information

GENERAL INFORMATION

SANMITE Miticide/Insecticide is intended for control of mites and whiteflies on ornamental plants, flowers, and foliage crops. SANMITE provides excellent knockdown and residual control. A good evaluation of performance can generally be made 4-7 days after treatment.

Crop Tolerance

All crops listed. Plant Species Tested for Tolerance to SANMITE are tolerant to SANMITE.

Mode of Action

SANMITE works primarily through contact action. Treat plants when pests are immature or at a susceptible stage and populations are building, before crop damage occurs.

Resistance Management

Using SANMITE in successive miticide applications is not recommended. Use SANMITE as part of a sound resistance management program that includes rotation with other treatments having different modes of action.

Spray Coverage

Apply SANMITE in sufficient water to ensure thorough coverage of foliage. Thorough coverage is required for optimum control. To achieve adequate coverage, use proper spray pressure, nozzles, nozzle spacing, and volume per acre. Consult spray nozzle and accessory guide for information pertaining to proper equipment calibration.

GENERAL RESTRICTIONS AND LIMITATIONS - ALL CROPS

- Do not enter a treated greenhouse or a treated indoor area without protective

equipment for 12 hours unless one of the following items is completed:

- 10 air exchanges
- 2 hours of system ventilation
- 4 hours of ventilation using vents, windows or other passive ventilation
- All required PPE is worn.
- Sequential Treatment: Do not use SANMITE in successive miticide applications. Use SANMITE in rotation with other treatments having different modes of action.
- Do not apply this product through any type of irrigation system.
- Do not apply this product aerially.
- Do not use SANMITE with nutritional sprays that contain boron.
- Do not apply this product as a smoke, mist, fog, or aerosol.
- Do not repackage or reformulate without manufacturer's written approval. For end use only.

Method

[Spray](#)

Rates

[field rates 0](#)

[field rates 1](#)

•

Restricted Entry Interval

12 hours

Timings

[When pests are immature or at a susceptible stage and populations building, before crop damage occurs.](#)