

LAWN AND PERIMETER - WEEVIL, BILLBUGS, ETC.

General Information

DIRECTIONS FOR USE

Not for use on plants being grown for sale of other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climate modification and being grown in interior plantscapes, ornamental gardens or parks, or lawns or grounds.

Do not apply more than 0.2 lb. active ingredient (a.i.) per acre per application on residential use sites (i.e., around private homes, apartment buildings, condominiums, non-agricultural outbuildings, non-commercial greenhouses, pre-schools or day care facilities). May be applied at up to 0.4 lb. a.i. per acre per application on non-residential use sites (i.e., around institutional, public, commercial or industrial buildings; parks; recreational areas or athletic fields).

Do not broadcast apply more than 0.4 lbs. A.I. (200 lbs. LESCO CrossCheck EZ Granular Insecticide) per Acre per year.

Not for use on golf courses, sod farms, nurseries, in commercial greenhouses or on grass grown for seed.

Resistance: Some insects are known to develop resistance to products used repeatedly for control. Because the development of resistance cannot be predicted, the use of the product should conform to resistance management strategies established for the use area. Consult your local or state pest management authorities for details. If resistance to the product develops in your area, the product, or other products with a similar mode of action, may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and suspect that resistance is a reasonable cause, immediately consult your local company representative or pest management advisor for the best alternative method of control for your area.

Limitations, Restrictions, and Exceptions

LAWN AND PERIMETER APPLICATIONS

Lawn Application: Broadcast LESCO CrossCheck EZ Granular Insecticide with suitable application equipment to ensure uniform coverage over the treatment area.

APPLICATION RATES

The application rates listed will provide excellent control of the respective pests under typical conditions. However, at the discretion of the applicator, LESCO CrossCheck EZ Granular Insecticide may be applied at up to 0.4 lb AI/A to control each of the pests listed. However, for residential lawn use, do not apply more than 0.2 lbs. A.I. per Acre per application.

In New York State, the product may NOT be applied to any grass or turf area within 100 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch).

In New York State, do make a single repeat application of LESCO CrossCheck EZ Granular Insecticide if there are signs of renewed insect activity, but not sooner than two weeks after the first application.

Perimeter Application: Apply to ground area 5 to 10 feet wide around and adjacent to the structure.

Boundary Applications: Apply to soil areas, ornamental gardens, pavement crevices, fencelines, planter areas, vegetation areas and mulch areas

(Refer to label for Pounds of LESCO CrossCheck EZ Granular for Perimeter Applications Linear Distance (Feet) Around Structure)

COMMENTS

Annual Bluegrass Weevil (*Hyperodes*) adults: Applications should be timed to control adult weevils as they leave their overwintering sites and move into grass areas. This movement generally begins when Forsythia is in full bloom and concludes when flowering dogwood is in full bloom. Consult your State Cooperative Extension Service for more specific information regarding application timing.

Billbug adults: Applications should be made when adult billbugs are first observed during April and May. Degree day models have been developed to optimize application timing. Consult your State Cooperative Extension Service for information

specific to your region. In temperate regions, spring applications targeting billbug adults will also provide control of over-wintered chinch bugs.

Black Turfgrass Ataenius adults: Applications should be made during May and July to control the first and second generation of black turfgrass ataenius adults, respectively. The May application should be timed to coincide with the full bloom stage of Vanhoutte spiraea (*Spiraea vanhouttei*) and horse chestnut (*Aesculus hippocastanum*). The July application should be timed to coincide with the blooming of Rose of Sharon (*Hibiscus syriacus*).

Method

[Broadcast](#)

Rates

[field_rates 0](#)

[field_rates 1](#)

[field_rates 2](#)

•

Timings

[when adult billbugs are first observed during April and May.](#)

[During May and July to control the first and second generation of black turfgrass ataenius adults, respectively.](#)