

## **GRAPES - GRAPE LEAF FOLDER, OMNIVOROUS LEAF ROLLER, ETC.**

### General Information

### USAGE CAUTIONS

Do not use for dried fruit.

Application of materials containing sulfur should not be made just prior to or during periods of excessively high temperatures or within 10 days to 2 weeks of an oil spray.

Do not use in a manner or at a time other than in accordance with label directions because animals, plant or crop injury, illegal residues or other undesirable results may occur.

Do not apply to sulfur sensitive crops. Do not apply during high temperatures. Do not use within 2 weeks of an oil treatment, on citrus within 3 to 6 weeks. When grown for processing, consult the processor before applying sulfur.

### Limitations, Restrictions, and Exceptions

#### GRAPES - GRAPE LEAF FOLDER, OMNIVOROUS LEAF ROLLER, ETC.

- Start treatment as soon as possible after hatching and before larvae are protected by leaf folds.
- For Powdery Mildew control, the product may be substituted for a regularly scheduled (7 to 10 day) application of dusting sulfur. But the product is not intended for use to control Powdery Mildew other than in a scheduled program.

### Method

[N. A.](#)

### Rates

[field\\_rates 0](#)

•

### Restricted Entry Interval

24 hours

Timings

After hatching and before larvae are protected by leaf folds.