

FIELD CORN, ETC. - AT PLANT USE - 60 DAYS OF HARVEST FOR FORAGE

General Information

GENERAL INSTRUCTIONS

Use low rate under light to moderate infestation. Higher rates should be used under heavy insect pressure. The rate of application is variable according to insect pressure, timing of spray and field scouting. In New York State this product may not be applied within 100 feet (using ground equipment) to 300 feet (using aerial equipment) of coastal marshes or streams that drain into coastal marshes.

Hero EW Insecticide contains the pyrethroids zeta-cypermethrin and bifenthrin.

Maximum Usage When Applying Zeta-Cypermethrin and Cypermethrin Products to the Same Crop Within the Same Season.

Do not apply more than the maximum seasonal total for either zeta-cypermethrin or cypermethrin products when used alone; do not apply more than the combined maximum seasonal total for both products as outlined in the table of the label.

Limitations, Restrictions, and Exceptions

FIELD CORN

RESTRICTIONS

PHI: Do not apply within 30 days of harvest for grain

Maximum Amount per Application: Do not apply more than 11.2 ounces of product or 0.10 pound active ingredient per acre as an at-plant application.

Maximum Amount of Hero EW Insecticide allowed per Season: Do not apply more than 44.8 ounces of product or 0.40 pounds active ingredient per acre per season including at-plant plus foliar applications for this product.

Refer to the maximum usage tables when applying more than one product containing either zeta-cypermethrin or bifenthrin to this crop.

Use of this product on corn is prohibited in all coastal counties.

REMARKS

For Seed cord maggot, Root aphids White grubs and Wireworms: Apply in-furrow or in a 3 - 4 inch T-Band (band over the open furrow) at planting in a minimum of 2 - 7 gallons per acre.

For Armyworm spp. and Cutworm spp: Apply at planting on the soil surface in a 5 - 7 inch band in a minimum of 2 - 7 gallons per acre or broadcast in a minimum of 10 gallons per acre.

Use higher recommended dosage for increased residual pest control.

Method

[In Furrow](#)

[Band](#)

[Broadcast](#)

Pre-Harvest Interval

60 days

Rates

[field rates 0](#)

[field rates 1](#)

•

Restricted Entry Interval

12 hours

Timings

[At-Plant](#)