

DRY PEAS AND BEANS

General Information

GENERAL INSTRUCTIONS

Use low rate under light to moderate infestation. Higher rates should be used under heavy insect pressure. The rate of application is variable according to insect pressure, timing of spray and field scouting. In New York State this product may not be applied within 100 feet (using ground equipment) to 300 feet (using aerial equipment) of coastal marshes or streams that drain into coastal marshes.

Hero EW Insecticide contains the pyrethroids zetacypermethrin and bifenthrin.

Maximum Usage When Applying Zeta-Cypermethrin and Cypermethrin Products to the Same Crop Within the Same Season.

Do not apply more than the maximum seasonal total for either zetacypermethrin or cypermethrin products when used alone; do not apply more than the combined maximum seasonal total for both products as outlined in the table of the label.

Limitations, Restrictions, and Exceptions

DIRECTIONS FOR USE:

- Through coverage is essential to achieve control. Apply by ground or air using sufficient spray volume to obtain full coverage of foliage. Multiple applications may be required.

DRY PEAS AND BEANS

- Use a minimum spray volume of 10 gallons by ground and 2 gallon by air.

Restrictions:

Do not make applications less than 7 days apart.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

21 days

Rates

[field rates 0](#)

-

Restricted Entry Interval

12 hours

Timings

[N.A.](#)