

## **SWAMPS, MARSHES, ETC. - HEAVIER OR DENSER VEGETATIVE COVER**

General Information

### **DIRECTIONS FOR USE**

BVA 2 Mosquito Larvicide may be used to control mosquito larvae in swamps, marshes, floodwater areas, drainage areas, storm sewer catch basins, waste treatment facilities, settling ponds, ditches, temporary rain pools and other man-made depressions containing stagnant water pools, including, abandoned swimming pools, fountains and ponds.

Apply BVA 2 Mosquito Larvicide at rates of 1-5 gallons per acre depending on how dense the vegetation and weeds may be. This product may be applied by the following methods: handgun or handwand sprayer and by air.

BVA 2 Mosquito Larvicide has a water-white clear color and is practically odorless.

BVA 2 Mosquito Larvicide does not depend on life-cycle time for effectiveness. As it kills by suffocation, mosquitoes do not develop resistance. When applied as directed this product does not leave a dark oily film.

**Ground Spray (Handgun or Handwand) Applications** Ground applications must be in compliance with the specifications indicated in the label regarding Wind Speed, Temperature Inversions, and Droplet Size. Apply using a nozzle height of no more than 4 feet above the ground, water surface, or crop canopy.

### **Aerial Applications**

Aerial applications must be in compliance with the specifications indicated in the label regarding Wind Speed, Temperature Inversions, and Droplet Size.

**Release Height:** Do not release spray at a height greater than 10 feet above the ground, water surface, or the tops of plants that are to be sprayed.

Boom Length: The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter.

Swath Adjustment: When applications are made with a cross-wind, the swath will be displaced downwind. The applicator must compensate for this displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind. Leave at least one untreated swath at the downwind edge of the treated field.

Limitations, Restrictions, and Exceptions

## TREATMENT RATES

To penetrate heavier or denser vegetative cover, or if water is substantially polluted, use 2½ to 5 gallons per acre (7½ to 15 fl. oz. per 1000 sq. ft.).

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

[Hand-held](#)

Rates

[field rates 0](#)

[field rates 1](#)

•

Timings

[N.A.](#)