

VEGETABLES

General Information

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

This product is intended solely for use on agricultural crops grown for commercial or research purposes, and in State and/or Federal invasive mollusk eradication operations. This product may not be applied to residential sites (i.e., in yards, gardens, around homes and apartments) and other sites such as the exterior areas around schools and daycare facilities, and athletic fields, playgrounds, parks, recreation areas, etc.

Broadcast applications and applications of this product directed to plant parts are prohibited except to grass grown for seed, taro, and as specified for use in State and/or Federal invasive mollusk eradication operations.

RESTRICTIONS

- This product is intended solely for use on agricultural crops grown for commercial or research purposes, and in State and/or Federal invasive mollusk eradication operations.
- This product may not be applied to residential sites (i.e., in yards, gardens, around homes and apartments) and other sites such as the exterior areas around schools and daycare facilities, and athletic fields, playgrounds, parks, recreation areas, etc.
- Broadcast applications and applications of this product directed to plant parts are prohibited except to grass grown for seed, taro and as specified for use in State and/or Federal invasive mollusk eradication operations.
- Do not allow bait pellet to come in contact with plant parts.
- Use only as a baiting treatment to be applied directly to the soil area beneath

listed plants.

- Read entire label before using.
- Use as band treatment between rows.
- Do not apply as a wide-area broadcast.
- Grower is responsible for residues on crops.
- Do not apply to dry soil. Make applications following irrigation.
- Do not re-water for 40 hours after application.
- Apply any time of day, but evening applications are preferred.
- Do not use in or around Hanalei, Kauai (USFW Refuge)

Wilco Blue 4.0 Slug S. Snail Bait is a soil surface treatment for control of snails and slugs in citrus groves, berry crops, fistful vegetable crops, ornamental gardens, and seed grasses.

When applied as directed, Wilco Hue 4,0 Slug & Snail Bait will withstand rain and sprinkling and will offer protection to gardens for several weeks. This product may be injurious to Daylilies and Clematis. Do not use on these plants.

For use on ornamentals, artichokes, blueberries, caneberries, strawberries, lettuce, cole crops and other leafy greens, mint, corn, soybeans and tomatoes apply in a band, between rows and/or directed towards base of the plants, by hand or with commercial application equipment designed for banded application of granular products.

For use on citrus, apply in a band under tree canopy, by hand or with commercial application equipment designed for banded application of granular products.

Limitations, Restrictions, and Exceptions

Minimum days between retreatment: 14 days

Method

[Band](#)

Pre-Harvest Interval

0 day

Rates

[field_rates 0](#)

-

Restricted Entry Interval

12 hours

Timings

[Any time of day, but evening applications are preferred.](#)