

# **FOR USE ON CONTAINERIZED ANNUAL AND PERENNIAL FLOWERING AND FOLIAGE CROPS AND TROPICAL PLANTS**

## General Information

### GENERAL INFORMATION

Do not apply this product through any type of irrigation system.

FOR USE IN: COMMERCIAL GREENHOUSES, GLASSHOUSES (WHERE PLANTS ARE GROWN IN CONTAINERS), AND ON ACTIVELY GROWING STOCK PLANTS.

CONFIGURE increases lateral or basal branching and promotes flowering in certain ornamental crops. In addition, applications of CONFIGURE reduce the overall height of the plant resulting in more compact and marketable plants.

Plant response to CONFIGURE is strongly influenced by cultural and environmental variables. Growing media, water/fertilizer management, temperature, light, greenhouse composition, and other cultural practices may impact plant response. Plant size, container size and cultivar or variety can also influence treatment response. Although multiple cultivars have been tested for sensitivity to CONFIGURE and have demonstrated high levels of tolerance, it is impossible to insure an acceptable response in all cultivars. First time users of CONFIGURE should first conduct trials on a limited number of plants to determine crop response.

### APPLICATION INFORMATION

Apply CONFIGURE as a foliar spray using standard foliar application spray equipment. Make sure that sufficient volumes are used to thoroughly wet foliage. Spray uniformity is equally important. Uniformly apply 1-2 quarts of finished spray solution to 100 sq. ft. of area. A high quality wetting agent or spray adjuvant, approved for use on your crop, may be added to spray solutions according to the manufacturer's use instructions.

Refer in the label for CONFIGURE DILUTION GUIDE.

### Limitations, Restrictions, and Exceptions

APPLICATION INSTRUCTIONS FOR USE ON CONTAINERIZED ANNUAL AND PERENNIAL FLOWERING AND FOLIAGE CROPS AND TROPICAL PLANTS

- Apply CONFIGURE as a foliar spray when plants are fully established during active growth i.e., at least 3 to 4 weeks after potting.
- Apply CONFIGURE at rates ranging from 50 to 500 ppm.
- Apply in a uniform spray volume of 2 quarts per 100 square feet of area.
- First time users of CONFIGURE should treat a limited number of plants starting with 50 ppm on annuals and 100 ppm on perennials and tropical plants; observe for plant responses and make rate adjustments accordingly.
- Multiple applications of CONFIGURE at 7-10 day spray intervals may be necessary to achieve optimum results (i.e., increased lateral growth).

## Method

[Foliar spray](#)

## Rates

[field\\_rates 0](#)

- 

## Restricted Entry Interval

12 hours

## Timings

[When plants are fully established during active growth.](#)