

LAWN APPLICATION - ANNUAL BLUEGRASS WEEVIL, BILLBUGS, ETC.

General Information

DIRECTIONS FOR USE

Do not apply more than 0.2 lb. active ingredient (a.i.) per acre per application on residential use sites (i.e., around private homes, apartment buildings, condominiums, non-agricultural outbuildings, non-commercial greenhouses, pre-schools or day care facilities). Application at up to 0.4 lb. a.i. per acre per application is allowed on non-residential use sites (i.e., around institutional, public, commercial or industrial buildings; parks; recreational areas or athletic fields).

Do not broadcast apply more than 0.4 lbs. a.i. (200 lbs. UP-Star Gold Granular) per acre per year.

For ground application only. Do not apply by air.

Do not apply within 25 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish farm ponds.

Do not use on golf courses, sod farms, nurseries, in commercial greenhouses or on grass grown for seed.

Limitations, Restrictions, and Exceptions

LAWN APPLICATION

Broadcast UP-Star Gold Granular Insecticide with appropriate application equipment which ensures uniform coverage.

Instructions:

- Annual Bluegrass Weevil (*Hyperodes*) Adults: Time application to control adult weevils as they leave their over-wintering sites and move into grass areas. This movement usually starts when Forsythia is in full bloom and finishes when flowering dogwood is in full bloom. Consult your State Cooperative Extension Service for more specific information regarding application timing.

- Billbug Adults: Make first application when adult billbugs are first seen during April and May. In temperate regions, spring applications for billbug adults will also control over-wintered chinch bugs.

- Black Turfgrass *Ataenius* Adults: Apply in May and July to control the first and second generation of black turfgrass *ataenius* adults, respectively. Time the May application to coincide with the full bloom stage of Vanhoutte spiraea (*Spiraea vanhouttei*) and horse chestnut (*Aesulus hippocastanum*). Time the July application to match the blooming of Rose of Sharon (*Hibiscus syriacus*).

Method

[Broadcast](#)

Rates

[field rates 0](#)

[field rates 1](#)

[field rates 2](#)

•

Timings

[Billbug Adults: When adult billbugs are first seen during April and May.](#)

[Black Turfgrass *Ataenius*: In May and July.](#)