

## **INDOOR USE—DILUTED - ANIMAL QUARTER USE (BROWN DOG TICKS, ETC.)**

### General Information

#### DIRECTIONS FOR USE

Do not apply space spray while food processing is underway. Cover or remove all food processing surfaces. Thoroughly wash all food processing surfaces before reuse. After spraying in bakeries, meat packing plants, food processing plants, etc., all benches, shelving, equipment, etc. where exposed food will be handled must be washed with an effective cleaning compound followed by a potable water rinse to remove all traces of contamination. Food processing operations do not have to be stopped while applying a wet spray with care and in accordance with the directions and cautions above in those establishments which do NOT operate under Federal meat, poultry, shell egg grading and egg products inspection programs.

Do not apply directly to food, water, or feed supplements. Do not contaminate milk or milk handling equipment. Wash udder and teats of dairy animals before milking.

#### Limitations, Restrictions, and Exceptions

#### INDOOR USE—DILUTED

**ANIMAL QUARTER USE - DAIRY AND BEEF BARNS, POULTRY HOUSES, SWINE HOUSING AND OTHER LISTED AREAS:** For use in and around listed use areas to control accessible stages of: Brown Dog Ticks, Clover Mites, Cockroaches, Crickets, Deer Flies, Earwigs, Face Flies, Firebrats, Fleas, Flies, Horn Flies, Hornets, Mosquitoes, Silverfish, Small Flying Moths, Spiders, Stable Flies, Ticks, Wasps, Yellow Jackets, Fannia Flies and Darkling Beetles.

Apply as a space spray in conventional ULV cold aerosol generators or foggers adjusted to deliver aerosol size droplets. Whenever possible, space spray application should be made in the late evening when insects are at rest. Apply undiluted at the rate of 0.25 to 1.0 fluid ounces per 1,000 cubic feet of space above the animals. This concentrate may also be diluted with water at the rate of 1 part concentrate to 10 parts water and applied at the rate of 1 to 2 fluid ounces per 1,000 cubic feet of space above the animals. Direct spray toward the upper portions

of the enclosure, above the animals, filling the room with mist or fog. Vacate treated area and ventilate before re-occupying.

Method

[Directed](#)

Rates

[field\\_rates 0](#)

[field\\_rates 1](#)

[field\\_rates 2](#)

•

Timings

[Late evening when insects are at rest.](#)