

INDOOR USE—DILUTED - ANIMAL QUARTER USE (BED BUGS AND MITES)

General Information

DIRECTIONS FOR USE

Do not apply space spray while food processing is underway. Cover or remove all food processing surfaces. Thoroughly wash all food processing surfaces before reuse. After spraying in bakeries, meat packing plants, food processing plants, etc., all benches, shelving, equipment, etc. where exposed food will be handled must be washed with an effective cleaning compound followed by a potable water rinse to remove all traces of contamination. Food processing operations do not have to be stopped while applying a wet spray with care and in accordance with the directions and cautions above in those establishments which do NOT operate under Federal meat, poultry, shell egg grading and egg products inspection programs.

Do not apply directly to food, water, or feed supplements. Do not contaminate milk or milk handling equipment. Wash udder and teats of dairy animals before milking.

Limitations, Restrictions, and Exceptions

INDOOR USE—DILUTED

ANIMAL QUARTER USE - DAIRY AND BEEF BARNS, POULTRY HOUSES, SWINE HOUSING AND OTHER LISTED AREAS:

Apply as a space spray in conventional ULV cold aerosol generators or foggers adjusted to deliver aerosol size droplets. Whenever possible, space spray application should be made in the late evening when insects are at rest.

To control Bed Bugs and Mites in poultry houses, prepare a dilution of 1 part concentrate to 25 parts water and spray crevices of roost poles, cracks in walls, and cracks in nests where the Bed Bugs and Mites hide.

Method

[Directed](#)

Rates

[field_rates 0](#)

-

Timings

[Late evening when insects are at rest.](#)