

## **DORMANT APPLICATIONS FOR CONTROL OF MULTIFLORA ROSE - SPOT CONCENTRATE (5 FEET CANOPY DIAMETER)**

General Information

### **PRODUCT INFORMATION**

The following directions apply to all uses of this product. Additional precautions and restrictions will be found in each specific use section.

Do not treat irrigation ditches or water used for crop irrigation or domestic uses. Do not apply this product through any type of irrigation system.

Do not apply more than 1 lb a.i./A per application, and no more than 2 applications per year.

### **MIXING AND APPLICATION**

UNLESS OTHERWISE SPECIFIED UNDER THE INDIVIDUAL USE HEADINGS OF THE LABEL, THE FOLLOWING DIRECTIONS APPLY TO ALL CROP AND NON-CROP USES OF THIS PRODUCT. REFER TO INDIVIDUAL USE SECTIONS FOR ADDITIONAL PRECAUTIONS, RESTRICTIONS, APPLICATION RATES AND TIMING.

This product is a water-soluble formulation that can be applied using water or sprayable fluid fertilizer as the carrier. If a fluid fertilizer is to be used, a compatibility test (see COMPATIBILITY TEST) should be made prior to tank mixing.

Ground or Aerial application equipment which will give good spray coverage of weed foliage should be used. HOWEVER, DO NOT USE AERIAL APPLICATION EQUIPMENT IF SPRAY PARTICLES CAN BE CARRIED BY WIND INTO AREAS WHERE SENSITIVE CROPS OR PLANTS ARE GROWING OR WHEN TEMPERATURE INVERSIONS EXIST.

Apply 3 to 50 gallons of diluted spray per treated acre when using ground application equipment, or 1 to 10 gallons of diluted spray per treated acre (2 to 20 gallons of diluted spray per treated acre for preharvest uses) in a water-based carrier when using aerial application equipment. Use the higher level of the listed spray volumes when treating dense or tall vegetation. Use coarse sprays.

Select nozzles designed to produce minimal amounts of fine spray particles. Spray with nozzles as close to the weeds as is practical for good weed coverage.

To avoid uneven spray coverage, this product should not be applied during periods of gusty wind or when wind is in excess of 15 mph. Avoid disturbing (e.g., cultivating or mowing) treated areas for at least 7 days following application.

### Best Stewardship Practices

This product provides effective broadleaf weed and brush control when properly applied. Best stewardship practices in all mixing, loading, and application operations not only maximize weed control, but also protect ground and surface waters and minimize off-target movement.

This chemical is known to leach through soil into ground water under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

### Limitations, Restrictions, and Exceptions

#### DORMANT APPLICATIONS FOR CONTROL OF MULTIFLORA ROSE

This product can be applied when plants are dormant as an undiluted spot-concentrate directly to the soil or as a Lo-oil Basal Bark treatment using an oil-water emulsion solution.

SPOT-CONCENTRATE applications of this product should be applied directly to the soil as close as possible to the root crown but within 6 to 8 inches of the crown. On sloping terrain, application should be made to the uphill side of the crown. Do not make application when snow or water prevents applying this product directly to the soil. The use rate of this product is dependent on the canopy diameter of the multiflora rose. Examples: Use this product at 1/4, 1, 2-1/4 fluid ounces of product respectively, for 5, 10, or 15 feet canopy diameters. Do not exceed a total of 2

quarts of this product per acre per year.

Method

[Soil incorporation](#)

Rates

[field rates 0](#)

- 

Restricted Entry Interval

24 hours

Timings

[When plants are dormant.](#)