

## **PEANUT - CUTWORMS, LESSER CORNSTALK BORER, ETC. (AT-PLANT/POSTPLANT TREATMENT)**

### General Information

#### GENERAL INFORMATION

#### SMARTBOX INFORMATION

The SmartBox system should be properly calibrated by a person familiar with operation of the system to assure an accurate application rate of Lorsban 15G. Lorsban 15G must be applied with the SmartBox system. Please refer to the SmartBox operator's manual for detailed information on system operation and calibration procedures. Do not attempt to open or tamper with the SmartBox container. Do not attempt to repair the SmartBox container. If repairs are necessary, contact your local AMVAC sales representative or the manufacturer of the SmartBox system.

Refer to the label for the Application Rates/1000 Ft of Row and Equivalent/Acre at Different Row Spacings.

#### Limitations, Restrictions, and Exceptions

#### PEANUT

At Plant Preventive Treatment: Apply Lorsban 15G in a 6 to 12 inch band over the row behind the planter shoe and in front of the press wheel. Incorporate granules to a depth of 1 inch into the soil with tines drag chains, or other suitable equipment. If the 7.5 oz rate per 1000 feet of row is used at planting time, then another application of 7.5 oz per 1000 feet of row may be needed postplant to extend control.

- Band Rescue Treatment: Apply Lorsban 15G in a 10 to 18 inch band over the fruiting zone.

#### Pest-Specific Use Directions:

Suppression of white mold: Best suppression of white mold (southern blight) is

obtained by applying the maximum rate of 15 oz per 1000 ft of row. Irrigation or rain following application is needed to enhance treatment effectiveness for suppression of white mold. Under conditions of heavy white mold pressure, a suitable fungicide may also be required and must be applied separately.

#### Specific Use Restrictions:

- Do not apply more than 15 oz of Lorsban 15G per 1000 ft of row per crop season or apply more than 4 lb active chlorpyrifos per season.
- Do not make a second application of Lorsban 15G or other product containing chlorpyrifos within 10 days of the first application.
- Do not feed peanut forage or hay to meat or dairy animals.
- The combined total of preplant and postplant applications of Lorsban 4E and Lorsban 15G or other product containing chlorpyrifos must not exceed 4 lb of active ingredient per acre per crop season.

#### Method

##### [Band](#)

#### Pre-Harvest Interval

21 days

#### Rates

##### [field rates 0](#)

- 

#### Restricted Entry Interval

24 hours

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

#### Timings

##### [At-Plant](#)

##### [Postplant](#)