

TOBACCO - CUTWORMS, FIRE ANT (SUPPRESSION), ETC.

General Information

GENERAL INFORMATION

SMARTBOX INFORMATION

The SmartBox system should be properly calibrated by a person familiar with operation of the system to assure an accurate application rate of Lorsban 15G. Lorsban 15G must be applied with the SmartBox system. Please refer to the SmartBox operator's manual for detailed information on system operation and calibration procedures. Do not attempt to open or tamper with the SmartBox container. Do not attempt to repair the SmartBox container. If repairs are necessary, contact your local AMVAC sales representative or the manufacturer of the SmartBox system.

Refer to the label for the Application Rates/1000 Ft of Row and Equivalent/Acre at Different Row Spacings.

Limitations, Restrictions, and Exceptions

TOBACCO

Preplant Broadcast Treatment: One week before transplanting, apply Lorsban 15G in combination with a rotary granular applicator, such as a fertilizer spreader, to evenly distribute the granules over the soil surface. Immediately following application, incorporate the granules uniformly into the soil to a depth of 2 to 4 inches using a rotary hoe or other incorporation device. The application of Lorsban 15G will also suppress movement of imported fire ant into a treated field.

Specific Use Restrictions:

- Do not make more than 1 application of Lorsban 15G or other product containing chlorpyrifos per season.
- The maximum single application rate is 2.025 lb active chlorpyrifos per acre.

Method

Broadcast

Pre-Harvest Interval

7 days

Rates

field rates 0

-

Restricted Entry Interval

24 hours

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

Timings

Preplant