

TREE AND VINE CROPS - SOIL BROADCAST SPRAY INCORPORATED, FLOOD OR FURROW IRRIGATED

General Information

DIRECTIONS FOR USE

Designed to be applied to the soil.

FERTIL HUMUS should be applied when soil analysis identifies the need and proper rate of application for this product.

Best results will be obtained when application is concentrated in the active root zone. For tree & vine crops, first application should be 15 to 20 days before planting, bud break or root flush or at the onset of dormancy. For field crops, apply post harvest onto crop stubble, compost, or manure followed by shallow cultivation.

Can be applied in combination with compatible plant growth regulators, pesticides, or other liquid fertilizers.

If compatibility is in question, jar test a small quantity.

Do not apply this product in concentrations greater than 10%.

Do not apply HUMA GRO nutrients during the hottest time of the day or to plants under drought or flood stress.

Application timing, intervals and rates may vary according to individual crop requirement, stage of development, available nutrient levels in the soil, and overall nutritional status of the crop.

Applications can be made as often as every 15 to 20 days as needed.

Soil moisture is required for maximum bioactivity.

If soil is dry, moisture should be provided by irrigation.

Consult your local HUMA GRO Representative or other agricultural specialist for crop-specific recommendations.

Limitations, Restrictions, and Exceptions

TREE AND VINE CROPS

Tree or Vine: up to 3 quarts

Method

[In Furrow](#)

[Broadcast](#)

[Spray](#)

[Soil application](#)

[In Furrow](#)

[Broadcast](#)

[Spray](#)

[Soil application](#)

Rates

[field_rates 0](#)

•

Timings

[Preplant](#)

[At bud break or root flush or at the onset of dormancy.](#)