

CUCURBITS

General Information

DIRECTIONS FOR USE

Shake well before use. Avoid freezing.

For residential use in home gardens, lawns and ornamentals. Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.

MONTEREY GARDEN INSECT SPRAY is intended for control of worms (caterpillars) and other listed insects. MONTEREY GARDEN INSECT SPRAY does not significantly impact predatory beneficial insects, predatory mites, and spiders while controlling target pests. Susceptible insect pests may be observed on plants up to several hours after treatment, but will have ceased active feeding before being killed.

HOW TO MIX

Add the required amount of MONTEREY GARDEN INSECT SPRAY to the specified amount of water, mix thoroughly, and apply uniformly to both upper and lower surfaces of plant foliage. Mix only as much spray as needed for a single treatment. In vegetable gardens, for best results, do not use more than 3 gallons of spray for 1,000 sq. ft. of area.

Do not use kitchen utensils for measuring. Keep measuring utensils with product and away from children.

HOW TO APPLY (Shake well before use.)

MONTEREY GARDEN INSECT SPRAY may be applied with trigger sprayer, hand-held, backpack, or hose-end sprayers. Use a hose-end sprayer that can be adjusted to provide a dilution ratio of about 2 fl. oz. (4 Tbs.) of MONTEREY GARDEN INSECT SPRAY per gallon of spray. See Hose-end Sprayer Directions below if this product is packaged in hose-end sprayer.

WHEN TO APPLY

Apply when listed pests are present. Repeat applications may be made as indicated in the Home Gardens section. See your state extension service recommendations for treatment guidelines in your area.

USES

Home Gardens

In the state of Georgia, do not apply MONTEREY GARDEN INSECT SPRAY TO: Broccoli Raab, Chinese Cabbage (Bok Choy), Collards, Kale, Mizuna, Mustard Greens, Mustard Spinach, Rape Greens.

Limitations, Restrictions, and Exceptions

CUCURBITS

- Minimum Days To Wait Before Reapplying - 5
- Minimum Days To Wait From Last Application To Harvest (All except cucumber, 3 cucumber, 1)

Amount of Monterey Garden Insect Spray To Use per Pint, Quart or Gallon of Spray

- Per Pint (16 fl oz) of Spray
- Per Quart (32 fl oz) of Spray
- Per Gallon (128 fl oz) of Spray

Conversion factors: 1 fl oz = 30 ml = 2 tablespoons (Tbs) = 6 teaspoons (tsp)

(1 teaspoon = 1/3 tablespoon)

Method

[Spray](#)

Rates

[field_rates 0](#)

[field_rates 1](#)

[field_rates 2](#)

[field_rates 3](#)

[field_rates 4](#)

[field_rates 5](#)

-

Timings

[When listed pests are present.](#)