

# **MICE**

## General Information

### DIRECTIONS FOR USE

**IMPORTANT:** Do not expose children, pets or other nontarget animals to rodenticides.

To help prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of the reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hooved livestock, raccoons, bears, or other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container and unused, spoiled and unconsumed bait as specified on the label. Note: Bait stations are mandatory for outdoor, above-ground use. Tamper-resistant bait stations must be used if children, pets, non-target mammals or birds may access the bait.

**USE SITES:** The product can be used only to control Norway Rats, Roof Rats, and House Mice in and within 100 feet of man-made structures constructed in a manner so as to be vulnerable to commensal rodent invasions and/or to harboring or attracting rodent infestations. Examples of such structures include homes, and other permanent or temporary residences, food processing facilities, industrial and commercial buildings, trash receptacles, agricultural and public buildings, transport vehicles (ships, trains, aircraft), docks, and port of terminal buildings and related structures around and associated with these sites. Fence and perimeter baiting, beyond 100 feet from a structure as defined above is prohibited. The product must not be applied directly to food or feed crops.

USE RESTRICTIONS: Do not place near or inside ventilation duct openings. Do not contaminate water, food, feedstuffs, food or feed handling equipment, or milk or meat handling equipment or surfaces that come into direct contact with food. Do not use the product in sewers. Do not broadcast bait. Do not apply further than 100 feet from buildings. Do not sell the product in individual containers holding less than 4 pounds of bait.

BURROW BAITING WITH TOP GUN PLACE PACK IS PROHIBITED.

SELECTION OF TREATMENT AREAS: Determine areas where rats and mice will most likely find and consume the bait. Generally these are along wall, by gnawed openings, in or beside burrows, in comes and concealed place, between floors and walls, or in location where rats and mice or their signs have been seen. Remove as much alternative food accessible to rodents as possible.

APPLICATION DIRECTIONS: The active ingredient in the product is different from anticoagulation rodenticides. Individual rats and mice cease feeding after consuming a toxic dose.

FOLLOW-UP: Adjust the amount of bait applied to the level of rat and mouse feeding expected at each bait placement location. Highest bait consumption is expected to occur on the first day or two after treatment. After several days, inspect bait placements and replenish bait at sites where there is heavy feeding or where there is continued evidence of rat and mouse activity. Maintain bait for at least one week and continue baiting until all signs of rat and mouse activity have ceased or it becomes clear that the remaining rodents are not attracted to the bait.

Limitations, Restrictions, and Exceptions

MICE

Apply 1 to 2 place packs per placement. Space placements at intervals of 8 to 12 feet in infested areas. Larger placements, up to 6 place packs, may be needed at points of extremely high mouse activity.

Method

[Bait](#)

Rates

[field\\_rates 0](#)

- 

Timings

[N.A.](#)