

# **CONTROL OF MICE**

## General Information

**IMPORTANT:** Do not expose children, pets, or nontarget animals to rodenticides. To help to prevent accidents:

1. Store unused product out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from bait stations when they are lifted, units must be secured or otherwise immobilized. Stronger bait stations are needed in areas open to hooved livestock, raccoons, bears, or other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container and unused, spoiled, or unconsumed bait as specified on the label.

**USE RESTRICTIONS:** For control of Norway rats, roof rats and house mice in and around homes, industrial, commercial, agricultural and public buildings, and similar manmade structures. May also be used in transport vehicles (ships, trains, aircraft) and in and around related port or terminal buildings. This product may also be used in alleys. Do not use in sewers. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast bait. When used in USDA inspected facilities, this product must be applied in tamper-resistant bait stations.

**SELECTION OF TREATMENT AREAS:** Determine areas where rats or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been seen. Protect bait from rain and snow. Remove as much alternative food as possible.

## Limitations, Restrictions, and Exceptions

## APPLICATION DIRECTIONS:

(Qualifier statement for 10 x 0.88 oz. packages only: This package contains enough bait (8.8 oz.) to kill two to four rats).

MICE: Apply 1 pac per placement. Space placements at 8 to 12 foot intervals. Two pacs may be applied at points of extremely high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days or until there no longer are signs of fresh mouse activity.

FOLLOW-UP: Replace contaminated or spoiled bait immediately. Using gloves, collect and dispose of all dead, exposed animals and leftover bait. To prevent reinfestation, limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish as needed.

Method

[Bait](#)

Rates

[field\\_rates 0](#)

•

Timings

[N. A.](#)