

TURFGRASS - FAIRY RING

General Information

PRODUCT INFORMATION

Mixing Procedures

Prepare no more spray mixture than is needed for the immediate operation. Thoroughly clean spray equipment before using this product. Agitation is necessary for proper dispersal of the product. Maintain agitation throughout the spraying operation. Do not let the spray mixture stand overnight in the spray tank. Flush the spray equipment thoroughly following each use and apply the rinsate to a previously treated area.

Alone: Add 1/2 of the required amount of water to the mix tank. With the agitator running, add the TARTAN FUNGICIDE to the tank. Continue agitation while adding the remainder of the water. Begin application of the solution after TARTAN FUNGICIDE has completely dispersed into the mix water. Maintain agitation until all of the mixture has been applied.

TARTAN FUNGICIDE + Tank Mixtures: Add 1/2 of the required amount of water to the mix tank. Start the agitator running before adding any tank mix partners. In general, tank mix partners should be added in this order: (1) products packaged in water-soluble packaging, wettable powders, wettable granules (dry flowables); (2) liquid flowables, liquids; and (3) emulsifiable concentrates. Always allow each tank mix partner to become fully dispersed before adding the next product. Provide sufficient agitation while adding the remainder of the water. Maintain agitation until all of the mixture has been applied.

Note: When using TARTAN FUNGICIDE in tank mixtures, all products in water-soluble packaging should be added to the tank before any other tank mix partner. Allow the water-soluble packaging to completely dissolve and the product(s) to completely disperse before adding any other tank mix partner to the tank.

If using TARTAN FUNGICIDE in a tank mixture, observe all directions for use, crop/sites, use rates, dilution ratios, precautions, and limitations that appear on the tank mix product label. No label dosage rate should be exceeded, and the most restrictive label precautions and limitations should be followed. This product should

not be mixed with any product that prohibits such mixing. Tank mixtures or other applications of products referenced on this label are permitted only in those states in which the referenced products are labeled.

TARTAN FUNGICIDE is compatible with most insecticide, fungicide, and foliar nutrient products. However, the compatibility of TARTAN FUNGICIDE with tank mix partners should be tested before use.

To determine biological compatibility with other products, mix the products in the desired proportions, spray on target plants and observe for phytotoxicity seven days after the application.

To determine the physical compatibility of TARTAN FUNGICIDE with other products, use a jar test, as described below. Using a quart jar, add the proportionate amounts of the products to 1 qt of water. Add wettable powders and water-dispersible granular products first, then liquid flowables, and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank.

Observe all directions, precautions, and limitations on labeling of all products used in tank mixes. Tank mixtures or other applications of products referenced on this label are permitted only in those states in which the referenced products are registered.

Use with additives: Use of spray additives is not required. Any spray additive should be evaluated prior to use. Do not use in conjunction with organosilicate-based products, or plant injury may occur. Label directions are based on data with no additives.

Chemigation: Do not apply this product through any type of irrigation system.

Aerial Application: Do not apply by aerial application in the State of New York. Aerial application to turfgrass is limited to sodfarm turf only.

Resistance Management: TARTAN FUNGICIDE contains the strobilurin class of chemistry, which exhibits no known cross-resistance to other chemical classes including sterol inhibitors, dicarboximides, benzimidazoles, anilinopyrimidines, or

phenylamides. However, certain fungal pathogens are known to develop resistance to products used repeatedly. Because resistance development cannot be predicted, the use of this product should conform to resistance management strategies. Such strategies may include rotating and/or tank mixing with products having different modes of action; or limiting the total number of applications per season. Bayer encourages responsible product stewardship to ensure effective long-term control of the fungal diseases on this label. See specific recommendations in the turf section and the ornamentals section.

Maximum Use Rates

For turfgrass, up to 345 fl oz of TARTAN FUNGICIDE can be applied per acre per year. For turfgrass, maximum single application rate is 2.7 lb ai/A. (triadimefon).

Limitations, Restrictions, and Exceptions

TURFGRASS DISEASE CONTROL

TARTAN is a preventive and curative fungicide that may be applied to turf sites including golf courses, institutional and commercial lawns, sod farms, sports fields, parks, municipal grounds and cemeteries.

TARTAN is a broad spectrum fungicide for the control of dollar spot (*Sclerotinia homoeocarpa*), brown patch (*Rhizoctonia* spp), anthracnose (*Colletotrichum graminicola*), red thread (*Corticium fuciforme*), pink patch (*Limonomyces roseipellis*), Leaf spot (*Bipolaris* spp., *Drechslera* spp.), gray leaf spot (*Pyricularia grisea*), rust (*Puccinia* spp.), pink snowmold (*Microdochium nivale*), fusarium patch (*Microdochium nivale*), southern blight (*Sclerotium rolfsii*), stripe smut (*Ustilago striiformis*), and summer patch (*Magnaporthe poae*).

When conditions are favorable for Pythium blight (*Pythium* spp.), mix a Pythium control fungicide with TARTAN.

1. FUNGICIDES SUCH AS TARTAN ARE BEST USED IN A PREVENTIVE DISEASE CONTROL PROGRAM. CURATIVE APPLICATIONS CAN BE MADE FOR CERTAIN DISEASES.

2. It is recommended to apply in 1-2 gals. of water per 1,000 sq. ft. to ensure thorough coverage. For soil-borne diseases, use sufficient water to reach the crown and upper root zone.
3. Apply after mowing OR allow sprayed area to completely dry before mowing.
4. For control of both foliar and soil-borne diseases, allow sprayed area to completely dry before irrigation.
5. Under conditions optimum for high disease pressure, use the higher rate and the shorter interval.
6. For optimum turf quality and disease control, use TARTAN in conjunction with turf management practices that promote good health and optimum disease control.
7. Before use of any fungicide, proper diagnosis of the organism causing the disease is important. Use of diagnostic kits or other means of identification of the disease organism is essential to determine the best control measures.
8. Application to golf courses, including tees, greens, fairways, and roughs is permitted only if the turfgrass is 2.5 inches or less in height.

TURFGRASS

Preventive Application: The first application should be made winter/spring when 5-day average daily soil temperatures taken at 2 inches are 55 - 60°F. In sandy soils, average air temperature can also be an adequate measure. In areas, like southern Florida, where the soil temperatures do not get to 55-60°F, make the first application in mid to late January. Do not apply if day-time temperatures are above 80°F. Discontinue applications of plant growth regulators (PGR's) one week before and after the Tartan applications.

- Apply using 2 gallons of spray per 1,000 square feet.
- Applications should be watered in with ½ to ¾ inch of irrigation within 24 hours
Tank-mixing of wetting agents is not recommended with these preventive applications.
- A second application is required. Apply a second application 28 days later

Tank-mixing has resulted in a slight decrease in fungicide efficacy. Wetting agents should still be applied according to regular scheduling, preferably two weeks apart from the fungicide applications.

For preventive applications apply when conditions are favorable for disease. Use the higher rate/shorter interval for curative applications or under severe disease conditions.

Method

[Spray](#)

Rates

[field rates 0](#)

-

Restricted Entry Interval

24 hours

Exception: If the product is applied by drenching, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

Timings

[Winter/spring when 5-day average daily soil temperatures taken at 2 inches are 55 - 60°F.](#)