

## **COMBINATION DISINFECTION AND CLEANING (LIGHTLY TO HEAVILY SOIL)**

### General Information

#### DIRECTIONS FOR USE

SaniDate 5.0 Sanitizer/Disinfectant works best when diluted with water containing minimal levels of organic or inorganic materials, and with water having a neutral pH. Thoroughly rinse out tank with water before mixing concentrate. This product will readily mix with clean, neutral water and does not require agitation.

SaniDate 5.0 Sanitizer/Disinfectant concentrate should not be combined or mixed with any other pesticide concentrates.

This product is not for use on medical device surfaces.

Before use in federally inspected meat and poultry food processing plants and dairies, food products and packaging materials must be removed from room or carefully protected. A potable water rinse is not allowed (Do not rinse) following use of the product as a sanitizer on previously cleaned hard surfaces provided that the surfaces are adequately drained before contact with food so that little or no residue remains.

#### Limitations, Restrictions, and Exceptions

#### COMBINATION DISINFECTION AND CLEANING

Use a rate of 0.5 fl. oz. per gallon for hard, non-porous surfaces that are lightly soiled or have been pre-cleaned to remove gross contamination. For heavily soiled, hard, non-porous surfaces use a rate of 1 fl. oz per 1 gallon of water. Apply solution with mop, cloth, sponge, brush, scrubber, or coarse spray device or by soaking so as to wet all surfaces thoroughly. Allow surface to remain wet for ten (10) minutes then remove solution and entrapped soil with a clean wet mop, cloth, or wet vacuum pickup. Prepare a fresh solution daily or when it becomes soiled or diluted.

For treating sewer backups and for flooding remediation, prepare disinfecting solution of SaniDate 5.0 Sanitizer/Disinfectant by adding 1 fl. oz. of the product to 1 gallon of potable water. Remove gross filth from surfaces by cleaning with SaniDate

5.0 Sanitizer/Disinfectant solution by wiping, mopping, or as a coarse spray. Applications involving treatment of food contact surfaces, require a sterile or potable water rinse following cleaning.

Method

[Spray](#)

Rates

[field\\_rates 0](#)

•

Timings

[N.A.](#)